

Table of Contents

NMSSM spectrum calculators.....	1
(C++).....	1
NMSSMCALC (Fortran).....	1
NMSSMTools (Fortran77).....	1
(C++).....	1
SPheno (Fortran 90).....	1

NMSSM spectrum calculators

Email: Nikos Rompotis, Mario Pelliccioni, Ian Low, and Maggie Muehlleitner.

This is a list of NMSSM spectrum calculators that are currently under investigation, with a small summary of their main features:

(C++)

full 2-loop RGEs; full 1-loop corrections to all SUSY masses; 2-loop Higgs corrections $O(\alpha_s \alpha_t + \alpha_s \alpha_b + \alpha_{\tau}^2 + \alpha_t^2 + \alpha_t \alpha_b + \alpha_b^2)$; Z3-invariant and Z3-violating NMSSM; boundary conditions and EWSB output can be customized via a modelfile

NMSSMCALC (Fortran)

Real and complex NMSSM: Higgs boson masses up to $O(\alpha_t \alpha_s)$, branching ratios including the state-of-the-art HO (QCD, SUSY-QCD&EW - Δ_b ?, $\Delta_s \Delta_{\tau}$; , SUSY-QCD to $\sim q$ decays) corrections and off-shell decays.

NMSSMTools (Fortran77)

2-loop RGEs; computation of the Higgs (up to $O((\alpha_b + \alpha_t)\alpha_s)$) and sparticle masses and decay ratios, and the dark matter relic density via MicrOmegas. Possible variants are the general NMSSM, the Z3-invariant NMSSM, constraints at the GUT scale and GMSB. Present experimental constraints are implemented.

(C++)

full 2-loop RGEs; full 1-loop corrections to all SUSY masses; 2-loop Higgs corrections $O(\alpha_s \alpha_t + \alpha_s \alpha_b + \alpha_{\tau}^2 + \alpha_t^2 + \alpha_t \alpha_b + \alpha_b^2)$; Z3-invariant and Z3-violating NMSSM; a number of boundary conditions (mSUGRA, non-universal trilinears, general, ...)

SPheno (Fortran 90)

2-loop RGEs with flavour and CP effects; 1-loop corrections to all SUSY masses; 2-loop Higgs corrections beyond $O(\alpha_s(\alpha_b + \alpha_t))$; independent calculation of decays as well as flavour and precision observables.

Definition: $\alpha_{t,b} = h_{t,b}^2 / (4\pi)$, with $h_{t,b}$ being the superpotential top and bottom couplings. The one-loop $O(\alpha_t)$ corrections are in fact corrections proportional to $h_t^2 m_t^2$, i.e. $h_t^4 v^2$.

-- EricFeng - 2018-02-28

This topic: LHCPhysics > NMSSMSpectrumCalculators

Topic revision: r3 - 2018-03-01 - EricFeng



Copyright &© 2008-2021 by the contributing authors. All material on this collaboration platform is the property of the contributing authors.

or Ideas, requests, problems regarding TWiki? use Discourse or Send feedback