

method	baseline		baseline & $\cancel{H}_T > 250 \text{ GeV}$		baseline & $H_T > 500 \text{ GeV}$	
	$Z \rightarrow \nu\bar{\nu}$ from γ +jets	26.2	± 4.8	7.1	± 2.2	8.4
$Z \rightarrow \nu\bar{\nu}$ from $W \rightarrow \ell\nu$ (combined)	17.2	± 7.9	6.3	± 3.8	3.8	± 3.2
$Z \rightarrow \nu\bar{\nu}$ from $Z \rightarrow \ell\bar{\ell}$ (combined)	16.8	$^{+13.2}_{-9.9}$	–	–	–	–
$t\bar{t}/W \rightarrow e, \mu + X$ lost-lepton method	33.0	± 8.1	4.8	± 1.8	10.9	± 3.4
$t\bar{t}/W \rightarrow \tau_{\text{hadr}} + X$ method	22.8	± 4.1	6.7	± 2.1	8.5	± 2.5
QCD Rebalance & Smear method	29.7	± 15.2	0.16	± 0.10	16.0	± 7.9
QCD factorization method	25.2	± 13.4	0.4	± 0.3	17.3	± 9.4
Total data-driven background	111.7	± 18.3	18.8	± 3.5	43.8	± 9.2
Observed in 36 pb^{-1} of data	111		15		40	