

LHCHXSWG-INT-2015-001

March 30, 2015

LHC HIGGS CROSS SECTION WORKING GROUP\*

INTERNAL NOTE

---

**Recommendations for the interpretation of LHC  
searches for  $H_5^0$ ,  $H_5^\pm$ , and  $H_5^{\pm\pm}$  in vector boson  
fusion with decays to vector boson pairs**

---

Heather E. Logan<sup>1,†</sup> and Marco Zaro<sup>2,3,‡</sup>

<sup>1</sup> *Ottawa-Carleton Institute for Physics, Carleton University,  
1125 Colonel By Drive, Ottawa, Ontario K1S 5B6, Canada*

<sup>2</sup> *Sorbonne Universités, UPMC Univ. Paris 06, UMR 7589, LPTHE,  
F-75005, Paris, France*

<sup>3</sup> *CNRS, UMR 7589, LPTHE, F-75005, Paris, France*

---

\* <https://twiki.cern.ch/twiki/bin/view/LHCPhysics/LHCHXSWG>

† [logan@physics.carleton.ca](mailto:logan@physics.carleton.ca)

‡ [zaro@lpthe.jussieu.fr](mailto:zaro@lpthe.jussieu.fr)

## Abstract

We provide theory input for the interpretation of the LHC searches for the production of Higgs bosons  $H_5^0$ ,  $H_5^\pm$ , and  $H_5^{\pm\pm}$  that transform as a five-plet under the custodial symmetry. We choose as a benchmark the Georgi-Machacek model, in which isospin-triplet scalars are added to the Standard Model Higgs sector in such a way as to preserve custodial SU(2) symmetry. This leads to several interesting features, including a tree-level  $H_5^\pm W^\mp Z$  interaction,  $H_5^{\pm\pm}$  decays to like-sign  $W$  bosons, and  $H_5^0$  decays to  $W^+W^-$  and  $ZZ$  in a different ratio than appears in the Standard Model, while still satisfying experimental constraints on the electroweak  $\rho$  parameter. We provide cross sections for single production of all five scalars in vector boson fusion up to next-to-next-to-leading order in QCD for masses in the range 200–1000 GeV (200–2000 GeV) at the 8 TeV (13 TeV) LHC. We also provide tree-level decay widths for  $H_5^0 \rightarrow W^+W^-$  and  $ZZ$ ,  $H_5^\pm \rightarrow W^\pm Z$ , and  $H_5^{\pm\pm} \rightarrow W^\pm W^\pm$  for masses in the range 200–2000 GeV.