

References

Notes:

- a) CMS Notes are available at <http://cms.cern.ch/iCMS/> unless otherwise noted.
- b) References marked **doi** should be prefixed with <http://dx.doi.org/>.

-
- [1] T. L. S. Group, "The Large Hadron Collider Conceptual Design," *CERN-AC-95-05* (1995) [arXiv:hep-ph/0601012](https://arxiv.org/abs/hep-ph/0601012).
 - [2] CMS Collaboration, "The Compact Muon Solenoid Letter of Intent," *CERN/LHCC 1992-3* (1992). LHCC/I 1.
 - [3] CMS Collaboration, "The Compact Muon Solenoid Technical Proposal," *CERN/LHCC 94-38* (1994). LHCC/P1.
 - [4] **ATLAS and CMS** Collaboration, J. G. Branson et al., "High transverse momentum physics at the Large Hadron Collider: The ATLAS and CMS Collaborations," *Eur. Phys. J. direct* **C4** (2002) N1, [arXiv:hep-ph/0110021](https://arxiv.org/abs/hep-ph/0110021).
 - [5] **LHC/LC Study Group** Collaboration, G. Weiglein et al., "Physics interplay of the LHC and the ILC," [arXiv:hep-ph/0410364](https://arxiv.org/abs/hep-ph/0410364).
 - [6] N. Krasnikov and V. Matveev, "Search for new physics at LHC," *Phys. Usp.* **47** (2004) 643, [arXiv:hep-ph/0309200](https://arxiv.org/abs/hep-ph/0309200).
 - [7] CMS Collaboration, "The CMS Physics Technical Design Report, Volume 1," *CERN/LHCC 2006-001* (2006). CMS TDR 8.1.
 - [8] "OSCAR: CMS Simulation Package Home Page."
<http://cmsdoc.cern.ch/oscar>.
 - [9] **GEANT4** Collaboration, S. Agostinelli et al., "GEANT4: A simulation toolkit," *Nucl. Instrum. Meth.* **A506** (2003) 250–303.
 - [10] "ORCA: CMS Reconstruction Package." Site located at
<http://cmsdoc.cern.ch/orca>.
 - [11] **CMS** Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 2.6: Fast simulation," *CERN/LHCC 2006-001* (2006) 55.
 - [12] J. Pumplin et al., "New generation of parton distributions with uncertainties from global QCD analysis," *JHEP* **07** (2002) 012, [arXiv:hep-ph/0201195](https://arxiv.org/abs/hep-ph/0201195).

- [13] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, "MRST2001: Partons and alpha(s) from precise deep inelastic scattering and Tevatron jet data," *Eur. Phys. J.* **C23** (2002) 73, arXiv:hep-ph/0110215.
- [14] S. Alekhin et al., "HERA and the LHC - A workshop on the implications of HERA for LHC physics: Proceedings Part A," 2005.
- [15] C. Seez et al., "Photon decay modes of the intermediate mass Higgs," in *Proceedings of the Large Hadron Collider Workshop*, Jarlskog, G. and Rein, D., ed., volume II. CERN 90-10, Aachen, 4-9 October, 1990.
- [16] CMS Collaboration, "The Electromagnetic Calorimeter Technical Design Report," CERN/LHCC 97-033 (1997). CMS TDR 4, Addendum CERN/LHCC 2002-027.
- [17] A. Favara and M. Pieri, "Confidence Level Estimation and Analysis Optimization," *Preprint DFF 278/4/1997* (1997).
- [18] **L3 Collaboration** Collaboration, M. Acciarri et al., "Search for the Standard Model Higgs Boson in e+e- Interactions at $161 \text{ GeV} < \sqrt{s} < 172 \text{ GeV}$," *Phys. Lett. B* **411** (1997) 373.
- [19] **CMS Collaboration**, M. Pieri et al., "Inclusive search for the Higgs boson in the $H \rightarrow \gamma\gamma$ channel," *CMS Note AN 2006/112* (2006).
- [20] M. Spira, "QCD effects in Higgs physics," *Fortsch. Phys.* **46** (1998) 203–284, arXiv:hep-ph/9705337.
- [21] A. Djouadi, J. Kalinowski, and M. Spira, "HDECAY: a program for Higgs boson decays in the Standard Model and its supersymmetric extension," *Comput. Phys. Commun.* **108** (1998) 56–74. doi:10.1016/S0010-4655(97)00123-9.
- [22] **CMS Collaboration**, M. Dubinin et al., "The vector boson fusion production with $H \rightarrow \gamma\gamma$," *CMS Note 2006/097* (2006).
- [23] **CMS Collaboration**, M. Lethuillier, "Search for SM Higgs boson with W/Z+H, $H \rightarrow 2\gamma$ channel," *CMS Note 2006/110* (2006).
- [24] T. Sjostrand, L. Lonnblad, and S. Mrenna, "PYTHIA 6.2: Physics and manual," arXiv:hep-ph/0108264.
- [25] "CMSIM home page." Located at <http://cmsdoc.cern.ch/cmsim/cmsim.html>.
- [26] W. Giele et al., "The QCD/SM working group: Summary report," arXiv:hep-ph/0204316.
- [27] M. Dobbs et al., "The QCD/SM working group: Summary report," arXiv:hep-ph/0403100.
- [28] T. Binoth, "Two photon background for Higgs boson searches at the LHC," arXiv:hep-ph/0005194.
- [29] Z. Bern, L. J. Dixon, and C. Schmidt, "Isolating a light Higgs boson from the di-photon background at the LHC," *Phys. Rev.* **D66** (2002) 074018, arXiv:hep-ph/0206194.

- [30] T. Binoth, J. P. Guillet, E. Pilon, and M. Werlen, "A next-to-leading order study of photon pion and pion pair hadro-production in the light of the Higgs boson search at the LHC," *Eur. Phys. J. Direct* **C4** (2002). doi:10.1007/s1010502c0007.
- [31] CMS Collaboration, L. Agostino et al., "HLT Selection of Electrons and Photons," *CMS Note* **2006/078** (2006).
- [32] CMS Collaboration, E. Meschi et al., "Electron Reconstruction in the CMS Electromagnetic Calorimeter," *CMS Note* **2001/034** (2001).
- [33] CMS Collaboration, N. Marinelli, "Track Finding and Identification of Converted Photons," *CMS Note* **2006/005** (2006).
- [34] CMS Collaboration, V. Litvin et al., "The Rejection of Background to the $H \rightarrow \gamma\gamma$ Process Using Isolation Criteria Based on Information from the Calorimeter and Tracker," *CMS Note* **2002-030** (2002).
- [35] CMS Collaboration, M. Pieri et al., "Distinguishing Isolated Photos from Jets," *CMS Note* **2006/007** (2006).
- [36] The LEP Collaborations: ALEPH, DELPHI, L3 and OPAL, "Lower Bound for the Standard Model Higgs Boson Mass from Combining the Results of the Four LEP Experiments," *CERN EP* **98-046** (1998).
- [37] CMS Collaboration, S. Baffioni et al., "Discovery potential for the SM Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-e^+e^-$ decay channel," *CMS Note* **2006/115** (2006).
- [38] E. Barberio, B. van Eijk, and Z. Was, "PHOTOS: A Universal Monte Carlo for QED radiative corrections in decays," *Comput. Phys. Commun.* **66** (1991) 115–128.
- [39] E. Barberio and Z. Was, "PHOTOS: A Universal Monte Carlo for QED radiative corrections. Version 2.0," *Comput. Phys. Commun.* **79** (1994) 291–308. doi:10.1016/0010-4655(94)90074-4.
- [40] M. Spira, "HIGLU: A Program for the Calculation of the Total Higgs Production Cross Section at Hadron Colliders via Gluon Fusion including QCD Corrections," arXiv:hep-ph/9510347.
- [41] A. Djouadi, J. Kalinowski, and M. Spira, "HDECAY: A Program for Higgs Boson Decays in the Standard Model and its Supersymmetric Extension," *Comput. Phys. Commun.* **108** (1998) 56–74, arXiv:hep-ph/9704448.
- [42] C. Zecher et al., "Leptonic Signals from off-shell Z Boson Pairs at Hadron Colliders," arXiv:hep-ph/9404295.
- [43] CompHEP Collaboration, E. Boos et al., "CompHEP 4.4: Automatic computations from Lagrangians to events," *Nucl. Instrum. Meth.* **A534** (2004) 250–259, arXiv:hep-ph/0403113.
- [44] S. Slabospitsky and L. Sonnenschein, "TopReX generator (version 3.25): Short manual," *Comput. Phys. Commun.* **148** (2002) 87–102. doi:10.1016/S0010-4655(02)00471-X.
- [45] M. Beneke et al., "Top quark physics," arXiv:hep-ph/0003033.

- [46] CMS Collaboration, S. Baffioni, C. Charlot, F. Ferri, D. Futyan, P. Meridiani, I. Puljak, C. Rovelli, R. Salerno, and Y. Sirois, "Electron Reconstruction in CMS," *CMS Note* **2006/040** (2006).
- [47] S. Abdullin et al., "Study of PDF and QCD scale uncertainties in $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ events at the LHC," *CMS Note* **2006/068** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [48] P. Meridiani and R. Paramatti, "Use of $Z \rightarrow e^+e^-$ events for ECAL calibration," *CMS Note* **2006/039** (2006).
- [49] V. Bartsch and G. Quast, "Expected signal observability at future experiments," *CMS Note* **2005-004** (2006).
- [50] S. Bityukov and N. Krasnikov, "New physics discovery potential in future experiments," *Mod. Phys. Lett.* **A13** (1998) 3235–3249.
- [51] M. Aldaya, P. Arce, J. Caballero, B. de la Cruz, P. Garcia-Abia, J. Hernandez, M. Josa, and E. Ruiz, "Discovery potential and search strategy for the Standard Model Higgs boson in the $H \rightarrow ZZ^* \rightarrow 4\mu$ decay channel using a mass-independent analysis," *CMS Note* **2006/106** (2006).
- [52] V. Karimaki et al., "CMKIN v3 User's Guide," *CMS IN* **2004-016** (2004).
- [53] M. Spira, "'SM Higgs boson production cross sections at NLO' and 'SM Higgs boson BRs'." Available at <http://cmsdoc.cern.ch/~anikiten/cms-higgs/>, 2005.
- [54] Particle Data Group Collaboration, S. Eidelman et al., "Review of particle physics," *Phys. Lett.* **B592** (2004) 1. doi:10.1016/j.physletb.2004.06.001.
- [55] F. Maltoni, "Theoretical Issues and Aims at the Tevatron and LHC," in *Proceedings of the 1st Hadron Collider Physics Symposium (HCP 2005)*. Les Diablerets, Switzerland, July, 2005. Available at <http://hcp-2005.web.cern.ch/HCP-2005>.
- [56] J. Campbell, "W/Z + B anti-B / jets at NLO using the Monte Carlo MCFM," arXiv:hep-ph/0105226. Talk given at 36th Rencontres de Moriond on QCD and Hadronic Interactions, Les Arcs, France, 17-24 Mar 2001.
- [57] S. Abdullin et al., "Relative Contributions of t- and s-Channels to the $ZZ \rightarrow 4\mu$ Process," *CMS Note* **2006/057** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [58] P. Bartalini et al., "NLO vs. LO: kinematical differences for signal and background in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ analysis," *CMS Note* **2006/130** (2006).
- [59] D. Acosta et al., "Measuring Muon Reconstruction Efficiency from Data," *CMS Note* **2006/060** (2006).
- [60] A. Read, "Modified Frequentist Analysis of Search Results (The CLs Method)," *1st Workshop on Confidence Limits, CERN, Geneva* (2000). CERN-EP/2000-005.

- [61] LEP Working Group for Higgs boson searches Collaboration, R. Barate et al., "Search for the standard model Higgs boson at LEP," *Phys. Lett.* **B565** (2003) 61–75, arXiv:hep-ex/0306033.
- [62] S. Abdullin, A. Drozdetskiy, et al., "GARCON: Genetic Algorithm for Rectangular Cuts Optimization. User's manual for version 2.0," arXiv:hep-ph/0605143. See also <http://drozdets.home.cern.ch/drozdets/home/genetic/>.
- [63] S. Abdullin et al., "Sensitivity of the Muon Isolation Cut Efficiency to the Underlying Event Uncertainties," *CMS Note* **2006/033** (2006). See also hep-ph/0604120, Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report. April 2006.
- [64] M. Aldaya et al., "A method for determining the mass, cross-section and width of the Standard Model Higgs boson using the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ decay channel," *CMS Note* **2006/107** (2006).
- [65] V. Bartsch, "Simulation of Silicon Sensors and Study of the Higgs Decay,".
- [66] M. Dittmar and H. Dreiner, "How to find a Higgs boson with a mass between 155 GeV to to 180 GeV at the CERN LHC," *Phys. Rev. D* **55** (1997) 167–172, arXiv:hep-ph/9608317. doi:10.1103/PhysRevD.55.167.
- [67] M. Dittmar and H. Dreiner, "LHC Higgs Search with $l^+\nu l^-\bar{\nu}$ final states," *CMS Note* **1997/083** (1997).
- [68] T. Sjostrand et al., "High-energy-physics event generation with PYTHIA 6.1," *Comput. Phys. Commun.* **135** (2001) 238–259. doi:10.1016/S0010-4655(00)00236-8.
- [69] T. Binoth, M. Ciccolini, N. Kauer, and M. Kramer, "Gluon-induced W W background to Higgs boson searches at the LHC," *JHEP* **03** (2005) 065, arXiv:hep-ph/0503094.
- [70] G. Davatz et al., "Effective k-factors for $H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu$ at the LHC," *JHEP* **0405** (2004) 009, arXiv:hep-ph/0402218. doi:10.1088/1126-6708/2004/05/009.
- [71] S. Frixione and B. Webber, "Matching NLO QCD computations and parton shower simulations," *JHEP* **0206** (2002) 029, arXiv:hep-ph/0204244.
- [72] G. Davatz, M. Dittmar, and A. Giolo-Nicollerat, "Standard Model Higgs Discovery Potential of CMS in the $H \rightarrow WW^{(*)} \rightarrow l\nu l\nu$ Channel," *CMS Note* **2006/047** (2006).
- [73] J. M. Campbell and R. K. Ellis, "An update on vector boson pair production at hadron colliders," *Phys. Rev. D* **60** (1999) 113006. doi:10.1103/PhysRevD.60.113006.
- [74] J. Campbell and F. Tramontano, "Next-to-leading order corrections to Wt production and decay," *Nucl. Phys.* **B726** (2005) 109–130. doi:10.1016/j.nuclphysb.2005.08.015.
- [75] CMS Collaboration, "The TriDAS Project Technical Design Report, Volume 2: Data Acquisition and High-Level Trigger," *CERN/LHCC* **2002-26** (2002). CMS TDR 6.2.
- [76] V. Drollinger, U. Gasparini, E. Torassa, and M. Zanetti, "Physics study of the Higgs decay channel $H \rightarrow WW^{(*)} \rightarrow 2\mu 2\nu$," *CMS Note* **2006/055** (2006).

- [77] S. V. Chekanov, "Jet algorithms: a mini review," in *Proceedings of the 14th Topical Conference on Hadron Collider Physics (HCP 2002)*. Karlsruhe, Germany, Sep-Oct, 2002. arXiv:hep-ph/0211298.
- [78] S. Bityukov and N. Krasnikov, "Uncertainties and discovery potential in planned experiments," *CMS CR 2002/05* (2002).
- [79] G. Davatz, A. Giolo-Nicollerat, and M. Zanetti, "Systematics uncertainties of the top background in the $H \rightarrow WW$ channel," *CMS Note 2006/048* (2006).
- [80] F. Maltoni and T. Stelzer, "MadEvent: Automatic event generation with MadGraph," *JHEP 02* (2003) 027. doi:10.1088/1126-6708/2003/02/027.
- [81] N. Kauer, "Top background extrapolation for $H \rightarrow WW$ searches at the LHC," arXiv:hep-ph/0404045.
- [82] S. Lowette, J. D'Hondt, J. Heyninck, and P. Vanlaer, "Offline Calibration of b-Jet Identification Efficiencies," *CMS Note 2006/013* (2006).
- [83] V. Drollinger et al., "Modeling the production of W pairs at the LHC," *CMS Note 2005/024* (2005).
- [84] C. Anastasiou et al., "High precision QCD at hadron colliders: Electroweak gauge boson rapidity distribution at NNLO," *Phys. Rev. D 69* (2004).
- [85] I. Belotelov et al., "Simulation of misalignment scenarios for CMS tracking devices," *CMS Note 2006-008* (2006).
- [86] M. Cvetič and P. Langacker, "Implications of Abelian Extended Gauge Structures From String Models," *Phys. Rev. D 54* (1996) 3570–3579, arXiv:hep-ph/9511378.
- [87] M. Cvetič and P. Langacker, "New Gauge Bosons from String Models," *Mod. Phys. Lett. A 11* (1996) 1247–1262, arXiv:hep-ph/9602424.
- [88] A. Leike, "The phenomenology of extra neutral gauge bosons," *Phys. Rept. 317* (1999) 143–250, arXiv:hep-ph/9805494.
- [89] C. T. Hill and E. H. Simmons, "Strong dynamics and electroweak symmetry breaking," *Phys. Rept. 381* (2003) 235–402, arXiv:hep-ph/0203079.
- [90] T. Han, H. Logan, B. McElrath, and L.-T. Wang, "Phenomenology of the little Higgs model," *Phys. Rev. D 67* (2003) 095004. doi:10.1103/PhysRevD.67.095004.
- [91] M. Cvetič and S. Godfrey, "Discovery and identification of extra gauge bosons," in *Electroweak symmetry breaking and new physics at the TeV scale*, T. Barlow and S. Dawson, eds., pp. 383–415. World Scientific, 1996. arXiv:hep-ph/9504216.
- [92] S. Godfrey, "Update of discovery limits for extra neutral gauge bosons at hadron colliders," in *Proceedings of Snowmass 2001*, p. P344. Snowmass, Colorado, June–July, 2001. arXiv:hep-ph/0201093.
- [93] L. Randall and R. Sundrum, "A large mass hierarchy from a small extra dimension," *Phys. Rev. Lett. 83* (1999) 3370–3373. doi:10.1103/PhysRevLett.83.3370.

- [94] M. R. Whalley, D. Bourilkov, and R. C. Group, "The Les Houches accord PDFs (LHAPDF) and LHAGLUE," arXiv:hep-ph/0508110.
- [95] J. L. Rosner, "Off peak lepton asymmetries from new Z_s ," *Phys. Rev.* **D35** (1987) 2244.
- [96] J. L. Rosner, "Forward-Backward Asymmetries in Hadronically Produced Lepton Pairs," *Phys. Rev.* **D54** (1996) 1078–1082, arXiv:hep-ph/9512299.
- [97] R. N. Mohapatra, S. A. Goldstein, and D. Money, eds., "Unification and supersymmetry: The Frontiers of quark - lepton physics". Springer-Verlag, 2002. 3rd Ed.
- [98] I. Belotelov et al., "Influence of misalignment scenarios on muon reconstruction," *CMS Note* **2006-017** (2006).
- [99] R. Cousins, J. Mumford, and S. Valuev, "Detection of Z' gauge bosons in the dimuon decay mode in CMS," *CMS Note* **2005-002** (2005).
- [100] R. Cousins, J. Mumford, and V. Valuev, "Detection of Z' gauge bosons in the dimuon decay mode in CMS," *CMS CR* **2004/050** (2006).
- [101] V. Bartsch and G. Quast, "Expected signal observability at future experiments," *CMS Note* **2005-004** (2005).
- [102] S. Wilks, "The large-sample distribution of the likelihood ratio for testing composite hypotheses," *Annals of Math. Stat.* **9** (1938) 60.
- [103] U. Baur, O. Brein, W. Hollik, C. Schappacher, and D. Wackerroth, "Electroweak radiative corrections to neutral-current Drell- Yan processes at hadron colliders," *Phys. Rev.* **D65** (2002) 033007, arXiv:hep-ph/0108274.
- [104] V. Zykunov, "Weak radiative corrections to Drell-Yan process for large invariant mass of di-lepton pair," arXiv:hep-ph/0509315.
- [105] A. De Roeck and S. Slabospitsky, "Evaluation of the PDF Uncertainties (in CMS)." Available at <http://cmsdoc.cern.ch/cms/PRS/gentools/www/pdfuncert/uncert.html>.
- [106] G. Cerminara. Private communication.
- [107] P. Langacker, R. W. Robinett, and J. L. Rosner, "New heavy gauge bosons in pp and $p\bar{p}$ collisions," *Phys. Rev.* **D30** (1984) 1470.
- [108] J. L. Rosner, "Observability of charge asymmetries for lepton pairs produced in present collider experiments," *Phys. Lett.* **B221** (1989) 85.
- [109] T. P. Cheng and L. F. Li, "Gauge theory of elementary particle physics". Oxford University Press, 1988.
- [110] R. Cousins, J. Mumford, and V. Valuev, "Measurement of Forward-Backward Asymmetry of Simulated and Reconstructed $Z' \rightarrow \mu^+ \mu^-$ Events in CMS," *CMS Note* **2005-022** (2005).
- [111] M. Dittmar, "Neutral current interference in the TeV region: The experimental sensitivity at the LHC," *Phys. Rev.* **D55** (1997) 161–166, arXiv:hep-ex/9606002.

- [112] C.-E. Wulz, “ Z' at LHC,” in *Proceedings of the 1996 DPF/DPB Summer Study on New Directions in High-Energy Physics*. 1996. Available at <http://www.slac.stanford.edu/pubs/snowmass96>.
- [113] J. C. Collins and D. E. Soper, “Angular distribution of dileptons in high-energy hadron collisions,” *Phys. Rev.* **D16** (1977) 2219.
- [114] B. C. Allanach, K. Odagiri, M. A. Parker, and B. R. Webber, “Searching for narrow graviton resonances with the ATLAS detector at the Large Hadron Collider,” *JHEP* **09** (2000) 019, arXiv:hep-ph/0006114.
- [115] R. Cousins, J. Mumford, J. Tucker, and V. Valuev, “Spin discrimination of new heavy resonances at the LHC,” *JHEP* **11** (2005) 046.
doi:10.1088/1126-6708/2005/11/046.
- [116] I. Belotelov et al., “Search for Randall-Sandrum Graviton Decay into Muon Pairs,” *CMS Note* **2006/104** (2006).
- [117] S. Esen and R. M. Harris, “Jet Triggers and Dijet Mass,” *CMS Note* **2006/069** (2006).
- [118] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.3: Monte Carlo Corrections,” *CERN/LHCC* **2006-001** (2006) 409.
- [119] CDF Collaboration, F. Abe et al., “Search for new particles decaying to dijets at CDF,” *Phys. Rev.* **D55** (1997) 5263–5268. doi:10.1103/PhysRevD.55.R5263.
- [120] DØ Collaboration, V. M. Abazov et al., “Search for new particles in the two-jet decay channel with the DØ detector,” *Phys. Rev.* **D69** (2004) 111101, arXiv:hep-ex/0308033.
- [121] DØ Collaboration, B. Abbott et al., “The dijet mass spectrum and a search for quark compositeness in $\bar{p}p$ collisions at $\sqrt{s} = 1.8\text{--}TeV$,” *Phys. Rev. Lett.* **82** (1999) 2457–2462. doi:10.1103/PhysRevLett.82.2457.
- [122] E. Eichten, K. D. Lane, and M. E. Peskin, “New Tests for Quark and Lepton Substructure,” *Phys. Rev. Lett.* **50** (1983) 811–814.
doi:10.1103/PhysRevLett.50.811.
- [123] K. D. Lane, “Electroweak and flavor dynamics at hadron colliders,” arXiv:hep-ph/9605257.
- [124] CDF Collaboration, F. Abe et al., “Measurement of dijet angular distributions at CDF,” *Phys. Rev. Lett.* **77** (1996) 5336–5341. doi:10.1103/PhysRevLett.77.5336.
- [125] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.6.1: Data-driven calibration strategy,” *CERN/LHCC* **2006-001** (2006) 421.
- [126] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.6.2: Dijet Balancing,” *CERN/LHCC* **2006-001** (2006) 422.
- [127] CDF Collaboration, D. Acosta et al., “Measurement of the Lifetime Difference between B_s Mass Eigenstates,” *Phys. Rev. Lett.* **94** (2005) 101803, arXiv:hep-ex/0412057. doi:10.1103/PhysRevLett.94.101803.

- [128] **DØ** Collaboration, "Measurement of the Lifetime Difference in the B_s System," *DØ Conference note 5052* (2006).
- [129] A. Belkov and S. Shulga, "Studies of angular correlations in the decays $B_s^0 \rightarrow J/\psi\phi$ by using the SIMUB generator," *Comput. Phys. Commun.* **156** (2004) 221–240, arXiv:hep-ph/0310096. doi:10.1016/S0010-4655(03)00465-X.
- [130] A. S. Dighe, I. Dunietz, and R. Fleischer, "Extracting CKM phases and $B_s - \bar{B}_s$ mixing parameters from angular distributions of non-leptonic B decays," *Eur. Phys. J.* **C6** (1999) 647–662. doi:10.1007/s100529800954.
- [131] A. S. Dighe, I. Dunietz, H. J. Lipkin, and J. L. Rosner, "Angular distributions and lifetime differences in $B_s \rightarrow J/\psi\phi$ decays," *Phys. Lett.* **B369** (1996) 144–150. doi:10.1016/0370-2693(95)01523-X.
- [132] **CDF** Collaboration, F. Abe et al., "Production of J/ψ mesons from χ_c meson decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **79** (1997) 578–583. doi:10.1103/PhysRevLett.79.578.
- [133] B. Cano-Coloma and M. A. Sanchis-Lozano, "Charmonia production in hadron colliders and the extraction of colour-octet matrix elements," *Nucl. Phys.* **B508** (1997) 753–767, arXiv:hep-ph/9706270. doi:10.1016/S0550-3213(97)00660-3.
- [134] S. Cucciarelli, M. Konecki, D. Kotlinski, and T. Todorov, "Track reconstruction, primary vertex finding and seed generation with the Pixel Detector," *CMS Note 2006/026* (2006).
- [135] K. Prokofiev and T. Speer, "A kinematic fit and a decay chain reconstruction library," *CERN Yellow Report 2005-002* (2005). Proc. of the 2004 Conference for Computing in High-Energy and Nuclear Physics (CHEP 04), Interlaken, Switzerland, 2004, available at <http://indico.cern.ch>.
- [136] I. Dunietz, R. Fleischer, and U. Nierste, "In pursuit of new physics with B_s decays," *Phys. Rev.* **D63** (2001) 114015. doi:10.1103/PhysRevD.63.114015.
- [137] **BABAR** Collaboration, B. Aubert et al., "Ambiguity-free measurement of $\cos(2\beta)$: Time-integrated and time-dependent angular analyses of $B \rightarrow J/\psi K\pi$," *Phys. Rev.* **D71** (2005) 032005. doi:10.1103/PhysRevD.71.032005.
- [138] I. Dunietz, H. Quinn, A. Snyder, W. Toki, and H. J. Lipkin, "How to extract CP-violating asymmetries from angular correlations," *Phys. Rev.* **D43** (1991) 2193–2208. doi:10.1103/PhysRevD.43.2193.
- [139] P. Vanlaer et al., "Impact of CMS Silicon Tracker Misalignment on Track and Vertex Reconstruction," *CMS Note 2006/029* (2006).
- [140] T. Speer et al., "Vertex Fitting in the CMS Tracker," *CMS Note 2006/032* (2006).
- [141] S. Heinemeyer, W. Hollik, and G. Weiglein, "FeynHiggs: A program for the calculation of the masses of the neutral CP-even Higgs bosons in the MSSM," *Comput. Phys. Commun.* **124** (2000) 76–89, arXiv:hep-ph/9812320.

- [142] S. Heinemeyer, W. Hollik, and G. Weiglein, "The masses of the neutral CP-even Higgs bosons in the MSSM: Accurate analysis at the two-loop level," *Eur. Phys. J.* **C9** (1999) 343–366, arXiv:hep-ph/9812472.
- [143] G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, and G. Weiglein, "Towards high-precision predictions for the MSSM Higgs sector," *Eur. Phys. J.* **C28** (2003) 133–143, arXiv:hep-ph/0212020.
- [144] S. Lehti, "Study of bbZ as a benchmark for MSSM bbH ," *CMS Note 2006/099* (2006).
- [145] G. Bagliesi, S. Dutta, S. Gennai, A. Kalinowski, M. Konecki, D. Kotlinski, F. Moortgat, A. Nikitenko, W. L., and S. Wakefield, "Tau jet reconstruction and tagging at High Level Trigger and off-line," *CMS Note 2006/028* (2006).
- [146] S. Kunori, R. Kinnunen, and A. Nikitenko, "Missing Transverse Energy Measurement with Jet Energy Corrections," *CMS Note 2001/040* (2001).
- [147] H. Pi et al., "Measurement of Missing Transverse Energy with the CMS detector at the LHC," *CMS Note 2006/035* (2006).
- [148] A. Kalinowski and A. Nikitenko, "Measurement of the τ -tagging efficiency using the $Z \rightarrow \tau\tau \rightarrow \mu + \text{hadrons} + X$ events," *CMS Note 2006/074* (2006).
- [149] S. Gennai, A. Nikitenko, and L. Wendland, "Search for MSSM Heavy Neutral Higgs Boson in $\tau\tau \rightarrow \text{two Jet Decay Mode}$," *CMS Note 2006/126* (2006).
- [150] A. Rizzi, F. Palla, and G. Segneri, "Track impact parameter based b-tagging with CMS," *CMS Note 2006/019* (2006).
- [151] A. Kalinowski, M. Konecki, and D. Kotlinski, "Search for MSSM Heavy Neutral Higgs Boson in $\tau\tau \rightarrow \mu + \text{jet Decay Mode}$," *CMS Note 2006/105* (2006).
- [152] A. Kalinowski, M. Konecki, and D. Kotlinski, "Search for MSSM heavy neutral Higgs boson in $\tau + \tau \rightarrow \mu + \text{jet decay mode}$," *CMS Note 2006/105* (2006).
- [153] R. Kinnunen and S. Lehti, "Search for the Heavy Neutral MSSM Higgs Bosons with the $H/A \rightarrow \tau\tau \rightarrow \text{Electron plus Jet Decay Mode}$," *CMS Note 2006/075* (2006).
- [154] S. Jadach, Z. Was, R. Decker, and J. H. Kuhn, "The tau decay library TAUOLA: Version 2.4," *Comput. Phys. Commun.* **76** (1993) 361–380.
doi:10.1016/0010-4655(93)90061-G.
- [155] A.-S. Giolo-Nicollerat *CMS Note* (2006).
- [156] C. Weiser, "A Combined Secondary Vertex Based B-Tagging Algorithm in CMS," *CMS Note 2006/014* (2006).
- [157] S. Haywood et al., "Electroweak physics," arXiv:hep-ph/0003275. In 'Standard Model Physics (and more) at the LHC', Geneva, 1999.
- [158] D. Chakraborty, J. Konigsberg, and D. L. Rainwater, "Review of top quark physics," *Ann. Rev. Nucl. Part. Sci.* **53** (2003) 301–351, arXiv:hep-ph/0303092.

- [159] D. Benedetti, S. Cucciarelli, C. Hill, J. Incandela, S. Koay, C. Riccardi, A. Santocchia, A. Schmidt, P. Torre, and C. Weiser, "Search for $H \rightarrow bb$ in association with a tt pair at CMS," *CMS Note* **2006/119** (2006).
- [160] M. L. Mangano, M. Moretti, F. Piccinini, R. Pittau, and A. D. Polosa, "ALPGEN, a generator for hard multiparton processes in hadronic collisions," *JHEP* **07** (2003) 001, arXiv:hep-ph/0206293. doi:10.1088/1126-6708/2003/07/001.
- [161] W. Beenakker, S. Dittmaier, M. Kraemer, B. Pluemper, M. Spira, and P. Zerwas, "NLO QCD corrections to $t\bar{t}H$ production in hadron collisions," *Nucl. Phys.* **B653** (2003) 151–203, arXiv:hep-ph/0211352.
- [162] V. Roberfroid et al., "Validation and status of the HLT Steering Code and default menu," *CMS IN* **2006/001** (2006).
- [163] E. James, Y. Maravin, M. Mulders, and N. Neumeister, "Muon Identification in CMS," *CMS Note* **2006/010** (2006).
- [164] A. Heister, O. Kodolova, V. Konopliyanikov, S. Petrushanko, J. Rohlf, C. Tully, and A. Ulyanov, "Measurement of Jets with the CMS Detector at the LHC," *CMS Note* **2006/036** (2006).
- [165] A. Santocchia, "Optimization of Jet Reconstruction Settings and Parton-Level Correction for the $t\bar{t}H$ Channel," *CMS Note* **2006/059** (2006).
- [166] J. D'Hondt, S. Lowette, O. Buchmuller, S. Cucciarelli, F. Schilling, M. Spiropulu, S. Mehdiabadi, D. Benedetti, and L. Pape, "Fitting of Event Topologies with External Kinematic Constraints in CMS," *CMS Note* **2006/023** (2006).
- [167] M. Mangano. Private communication.
- [168] X.-N. Wang and M. Gyulassy, "HIJING: A Monte Carlo model for multiple jet production in $p p$, $p A$ and $A A$ collisions," *Phys. Rev.* **D44** (1991) 3501–3516.
- [169] M. Bedjidian et al., "Hard probes in heavy ion collisions at the LHC: Heavy flavour physics," arXiv:hep-ph/0311048.
- [170] K. Eskola, V. Kolhinen, and C. Salgado, "The scale dependent nuclear effects in parton distributions for practical applications," *Eur. Phys. J.* **C9** (1999) 61–68, arXiv:hep-ph/9807297.
- [171] O. Kodolova, M. Bedjidian, and S. Petrouchanko, "Dimuon reconstruction in heavy ion collisions using a detailed description of CMS geometry," *CMS Note* **1999-004** (1999).
- [172] M. Bedjidian and O. Kodolova, "Quarkonia measurements in heavy-ion collisions in CMS," *CMS Note* **2006/089** (2006).
- [173] G. Sterman and S. Weinberg, "Jets from Quantum Chromodynamics," *Phys. Rev. Lett.* **39** (1977) 1436.
- [174] JADE Collaboration, W. Bartel et al., "Experimental Studies on Multi-Jet Production in $e^+ e^-$ Annihilation at Petra Energies," *Z. Phys.* **C33** (1986) 23.

- [175] JADE Collaboration, S. Bethke et al., "Experimental Investigation of the Energy Dependence of the Strong Coupling Strength," *Phys. Lett.* **B213** (1988) 235.
- [176] S. Catani, Y. L. Dokshitzer, M. Olsson, G. Turnock, and B. R. Webber, "New clustering algorithm for multi-jet cross-sections in e+e- annihilation," *Phys. Lett.* **B269** (1991) 432–438.
- [177] CDF Run II Collaboration, A. Abulencia et al., "Measurement of the inclusive jet cross section in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ TeV using a cone-based jet algorithm," arXiv:hep-ex/0512020.
- [178] DØ Collaboration, R. Strohmer, "Inclusive jet cross-sections and dijet azimuthal decorrelations with DØ," *PoS HEP2005* (2006) 051, arXiv:hep-ex/0601016.
- [179] J. Butterworth, J. Couchman, B. Cox, and B. Waugh, "KtJet: A C++ implementation of the K(T) clustering algorithm," *Comput. Phys. Commun.* **153** (2003) 85–96, arXiv:hep-ph/0210022.
- [180] U. Baur, R. K. Ellis, and D. Zeppenfeld, eds., "QCD and Weak Boson Physics in Run II". FERMILAB, Batavia, IL, USA, 2000. Prepared for Physics at Run II: QCD and Weak Boson Physics Workshop: Final General Meeting, Batavia, Illinois, 4-6 Nov 1999.
- [181] S. D. Ellis, J. Huston, and M. Tonnesmann, "On building better cone jet algorithms," *eConf C010630* (2001) P513, arXiv:hep-ph/0111434.
- [182] M. Wobisch. Private communication.
- [183] T. Sjostrand, "High-energy physics event generation with PYTHIA 5.7 and JETSET 7.4," *Comput. Phys. Commun.* **82** (1994) 74–90.
- [184] Z. Nagy, "Three-jet cross sections in hadron hadron collisions at next-to-leading order," *Phys. Rev. Lett.* **88** (2002) 122003, arXiv:hep-ph/0110315.
- [185] T. Kluge, K. Rabbertz, and M. Wobisch, "fast NLO - fast pQCD calculations for hadron-induced processes." Available at <http://hepforge.cedar.ac.uk/fastnlo>. To be published.
- [186] S. Moretti, M. R. Nolten, and D. A. Ross, "Weak corrections and high E(T) jets at Tevatron," arXiv:hep-ph/0503152.
- [187] DØ Collaboration, V. M. Abazov et al., "The inclusive jet cross-section in p anti-p collisions at $s^{1/2} = 1.8$ -TeV using the k(T) algorithm," *Phys. Lett.* **B525** (2002) 211–218, arXiv:hep-ex/0109041.
- [188] DØ Collaboration, B. Abbott et al., "High- p_T jets in $p\bar{p}$ collisions at $\sqrt{s} = 630$ GeV and 1800 GeV," *Phys. Rev.* **D64** (2001) 032003. doi:10.1103/PhysRevD.64.032003.
- [189] CDF Collaboration, A. A. Affolder et al., "Charged jet evolution and the underlying event in proton anti-proton collisions at 1.8-TeV," *Phys. Rev.* **D65** (2002) 092002.
- [190] CDF Collaboration, D. Acosta et al., "The underlying event in hard interactions at the Tevatron anti-p p collider," *Phys. Rev.* **D70** (2004) 072002, arXiv:hep-ex/0404004.

- [191] T. Sjostrand and M. van Zijl, "A Multiple Interaction Model for the Event Structure in Hadron Collisions," *Phys. Rev.* **D36** (1987) 2019.
- [192] J. Butterworth, J. R. Forshaw, and M. Seymour, "Multiparton interactions in photoproduction at HERA," *Z. Phys.* **C72** (1996) 637–646, arXiv:hep-ph/9601371.
- [193] T. Gleisberg et al., "SHERPA 1.alpha, a proof-of-concept version," *JHEP* **02** (2004) 056, arXiv:hep-ph/0311263.
- [194] F. W. Bopp, R. Engel, and J. Ranft, "Rapidity gaps and the PHOJET Monte Carlo," arXiv:hep-ph/9803437.
- [195] G. Corcella et al., "HERWIG 6: An event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)," *JHEP* **01** (2001) 010, arXiv:hep-ph/0011363. doi:10.1088/1126-6708/2001/01/010.
- [196] C. M. Buttar, D. Clements, I. Dawson, and A. Moraes, "Simulations of minimum bias events and the underlying event, MC tuning and predictions for the LHC," *Acta Phys. Polon.* **B35** (2004) 433–441.
- [197] CDF Collaboration, R. Field, "Min-bias and the underlying event in Run 2 at CDF," *Acta Phys. Polon.* **B36** (2005) 167–178.
- [198] D. Acosta et al., "The Underlying Event at the LHC," *CMS Note* (2006), no. 2006/067.
- [199] CDF Collaboration, F. Abe et al., "Measurement of the Z -boson p_T distribution in $\bar{p}p$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **67** (1991) 2937–2941. doi:10.1103/PhysRevLett.67.2937.
- [200] P. Bartalini, R. Chierici, and A. De Roeck, "Guidelines for the estimation of theoretical uncertainties at the LHC," *CMS Note* **2005/013** (2005).
- [201] W. Adam, B. Mangano, T. Speer, and T. Todorov, "Track reconstruction in the CMS tracker," *CMS Note* **2006/041** (2006).
- [202] V. Andreev, D. Cline, and S. Otwinowski, "Inclusive b quark production," *CMS Note* **2006/120** (2006).
- [203] CDF Collaboration, F. Abe et al., "Measurement of the bottom quark production cross section using semileptonic decay electrons in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **71** (1993) 500. doi:10.1103/PhysRevLett.71.500.
- [204] CDF Collaboration, F. Abe et al., "Measurement of the B meson and b quark cross sections at $\sqrt{s} = 1.8$ TeV using the exclusive decay $B^0 \rightarrow J/\psi K^*(892)^0$," *Phys. Rev. D* **50** (1994) 4252. doi:10.1103/PhysRevD.50.4252.
- [205] CDF Collaboration, F. Abe et al., "Measurement of the B Meson Differential Cross Section $d\sigma/dp_T$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **75** (1995) 1451. doi:10.1103/PhysRevLett.75.1451.
- [206] CDF Collaboration, F. Abe et al., "Measurement of the B^+ total cross section and B^+ differential cross section $d\sigma/dp_T$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. D* **65** (2002) 052005. doi:10.1103/PhysRevD.65.052005.

- [207] CDF Collaboration, F. Abe et al., "Measurement of the ratio of b quark production cross sections in $\bar{p}p$ collisions at $\sqrt{s} = 630$ GeV and $\sqrt{s} = 1800$ GeV," *Phys. Rev. D* **66** (2002) 032002. doi:10.1103/PhysRevD.66.032002.
- [208] DØ Collaboration, S. Abachi et al., "Inclusive μ and b -quark production cross-sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **74** (1995) 3548. doi:10.1103/PhysRevLett.74.3548.
- [209] DØ Collaboration, B. Abbott et al., "The $b\bar{b}$ production cross section and angular correlations in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Lett.* **B487** (2000) 264. doi:10.1016/S0370-2693(00)00844-3.
- [210] DØ Collaboration, B. Abbott et al., "Cross section for b -Jet production in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **85** (2000) 5068. doi:10.1103/PhysRevLett.85.5068.
- [211] H1 Collaboration, C. Adloff et al., "Measurement of open beauty production at HERA," *Phys. Lett.* **B 467** (1999) 156. doi:10.1016/S0370-2693(99)01099-0.
- [212] H1 Collaboration, C. Adloff et al., "Erratum," *Phys. Lett.* **B 518** (2001) 331.
- [213] ZEUS Collaboration, J. Breitweg et al., "Measurement of open beauty production in photoproduction at HERA," *Eur. Phys. J.* **C18** (2001) 625–637. doi:10.1007/s100520100571.
- [214] H1 Collaboration, A. Aktas et al., "Measurement of F2(c anti-c) and F2(b anti-b) at high Q^2 using the H1 vertex detector at HERA," *Eur. Phys. J.* **C40** (2005) 349–359, arXiv:hep-ex/0411046.
- [215] ZEUS Collaboration, S. Chekanov et al., "Bottom photoproduction measured using decays into muons in dijet events in e p collisions at $s^{1/2} = 318$ GeV," *Phys. Rev. D* **70** (2004) 012008.
- [216] ZEUS Collaboration, S. Chekanov et al., "Measurement of beauty production in deep inelastic scattering at HERA," *Phys. Lett.* **B599** (2004) 173–189, arXiv:hep-ex/0405069.
- [217] H1 Collaboration, A. Aktas et al., "Measurement of beauty production at HERA using events with muons and jets," *Eur. Phys. J.* **C41** (2005) 453–467, arXiv:hep-ex/0502010.
- [218] L3 Collaboration, M. Acciarri et al., "Measurements of the cross sections for open charm and beauty production in $\gamma\gamma$ collisions at $\sqrt{s} = 189$ GeV - 202 GeV," *Phys. Lett.* **B503** (2001) 10. doi:10.1016/S0370-2693(01)00134-4.
- [219] L3 Collaboration, P. Achard et al., "Measurement of the cross section for open-beauty production in photon photon collisions at LEP," *Phys. Lett.* **B619** (2005) 71.
- [220] P. Nason, S. Dawson, and R. K. Ellis, "The Total Cross-Section for the Production of Heavy Quarks in Hadronic Collisions," *Nucl. Phys.* **B303** (1988) 607.
- [221] P. Nason, S. Dawson, and R. K. Ellis, "The One Particle Inclusive Differential Cross-Section for Heavy Quark Production in Hadronic Collisions," *Nucl. Phys.* **B327** (1989) 49–92.

- [222] W. Beenakker, W. Van Neerven, R. Meng, G. Schuler, and J. Smith, "QCD corrections to heavy quark production in hadron-hadron collisions," *Nucl. Phys.* **B351** (1991) 507–560. doi:10.1016/S0550-3213(05)80032-X.
- [223] M. Cacciari, S. Frixione, M. L. Mangano, P. Nason, and G. Ridolfi, "QCD analysis of first b cross section data at 1.96-TeV," *JHEP* **07** (2004) 033. See also included references. doi:10.1088/1126-6708/2004/07/033.
- [224] M. L. Mangano, "The saga of bottom production in proton antiproton collisions," *AIP Conf. Proc.* **753** (2005) 247–260, arXiv:hep-ph/0411020. See also included references.
- [225] S. Frixione, "Bottom production," arXiv:hep-ph/0408317.
- [226] S. Weinberg, "Unitarity Constraints on CP Nonconservation in Higgs Exchange," *Phys. Rev.* **D42** (1990) 860–866.
- [227] L. Lavoura, "Maximal CP violation via Higgs boson exchange," *Int. J. Mod. Phys.* **A8** (1993) 375–390.
- [228] E. Jenkins, M. E. Luke, A. V. Manohar, and M. J. Savage, "Semileptonic B_c decay and heavy quark spin symmetry," *Nucl. Phys.* **B390** (1993) 463–473, arXiv:hep-ph/9204238.
- [229] V. V. Kiselev, A. E. Kovalsky, and A. K. Likhoded, " B_c decays and lifetime in QCD sum rules," *Nucl. Phys.* **B585** (2000) 353–382. doi:10.1016/S0550-3213(00)00386-2.
- [230] X.-G. Wu, C.-H. Chang, Y.-Q. Chen, and Z.-Y. Fang, "The meson B_c annihilation to leptons and inclusive light hadrons," *Phys. Rev.* **D67** (2003) 094001. doi:10.1103/PhysRevD.67.094001.
- [231] C.-H. Chang and Y.-Q. Chen, "The Production of B_c or \bar{B}_c meson associated with two heavy quark jets in Z^0 boson decay," *Phys. Rev.* **D46** (1992) 3845–3855. doi:10.1103/PhysRevD.46.3845.
- [232] C.-H. Chang and Y.-Q. Chen, "Hadronic production of the B_c meson at TeV energies," *Phys. Rev. D* **48** (1993) 4086. doi:10.1103/PhysRevD.48.4086.
- [233] C.-H. Chang, Y.-Q. Chen, G.-P. Han, and H.-T. Jiang, "On hadronic production of the B_c meson," *Phys. Lett. B* **364** (1995) 78. doi:10.1016/0370-2693(95)01235-4.
- [234] C.-H. Chang, Y.-Q. Chen, and R. Oakes, "Comparative study of the hadronic production of B_c mesons," *Phys. Rev. D* **54** (1996) 4344. doi:10.1103/PhysRevD.54.4344.
- [235] C.-H. Chang and X.-G. Wu, "Uncertainties in estimating hadronic production of the meson B/c and comparisons between TEVATRON and LHC," *Eur. Phys. J.* **C38** (2004) 267–276. doi:10.1140/epjc/s2004-02015-0.
- [236] CDF Collaboration, F. Abe et al., "Observation of the B_c meson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **81** (1998) 2432–2437. doi:10.1103/PhysRevLett.81.2432.

- [237] CDF Collaboration, D. Acosta et al., "Evidence for the exclusive decay $B_c^\pm \rightarrow J/\psi\pi^\pm$ and measurement of the mass of the B_c meson," *Phys. Rev. Lett.* **96** (2006) 082002. doi:10.1103/PhysRevLett.96.082002.
- [238] A. V. Berezhnoi, V. V. Kiselev, A. K. Likhoded, and A. I. Onishchenko, "B/c meson at LHC," *Phys. Atom. Nucl.* **60** (1997) 1729–1740, arXiv:hep-ph/9703341.
- [239] I. P. Gouz, V. V. Kiselev, A. K. Likhoded, V. I. Romanovsky, and O. P. Yushchenko, "Prospects for the B_c studies at LHCb," *Phys. Atom. Nucl.* **67** (2004) 1559–1570. doi:10.1134/1.1788046.
- [240] C.-H. Chang and Y.-Q. Chen, "Decays of the B_c meson," *Phys. Rev.* **D49** (1994) 3399–3411. doi:10.1103/PhysRevD.49.3399.
- [241] W.-k. Kwong and J. L. Rosner, "Masses of new particles containing b quarks," *Phys. Rev.* **D44** (1991) 212–219. doi:10.1103/PhysRevD.44.212.
- [242] E. J. Eichten and C. Quigg, "Mesons with beauty and charm: Spectroscopy," *Phys. Rev.* **D49** (1994) 5845–5856, arXiv:hep-ph/9402210.
- [243] Y.-Q. Chen and Y.-P. Kuang, "Improved QCD motivated heavy quark potentials with explicit $\Lambda_{\overline{MS}}$ dependence," *Phys. Rev.* **D46** (1992) 1165–1171. doi:10.1103/PhysRevD.46.1165.
- [244] G. Chen, X. Meng, and J. Tao, "Feasibility to study the B_c meson at CMS," *CMS Note* **2006/118** (2006).
- [245] T. Sjostrand, L. Lonnblad, S. Mrenna, and P. Skands, "PYTHIA 6.3: Physics and manual," arXiv:hep-ph/0308153.
- [246] M. Grothe et al., "Triggering on Forward Physics," *CMS Note* **2006/054** (2006). Also available as TOTEM NOTE 2006-01.
- [247] CMS/TOTEM Collaboration, "CMS/TOTEM document on diffractive and forward physics, in preparation," 2006.
- [248] UA8 Collaboration, R. Bonino et al., "Evidence for Transverse Jets in High Mass Diffraction," *Phys. Lett.* **B211** (1988) 239.
- [249] M. Arneodo and M. Diehl, "Diffraction for non-believers," arXiv:hep-ph/0511047.
- [250] UA8 Collaboration, A. Brandt et al., "Measurements of single diffraction at $\sqrt{s} = 630$ GeV: Evidence for a non-linear $\alpha(t)$ of the pomeron," *Nucl. Phys.* **B514** (1998) 3–44. doi:10.1016/S0550-3213(97)00813-4.
- [251] S. Erhan and P. E. Schlein, "Inelastic diffraction data and the Pomeron trajectory," *Phys. Lett.* **B481** (2000) 177–186. doi:10.1016/S0370-2693(00)00467-6.
- [252] K. Goulianos, "Twenty years of diffraction at the Tevatron," arXiv:hep-ph/0510035. Presented at 11th International Conference on Elastic and Diffractive Scattering: Towards High Energy Frontiers: The 20th Anniversary of the Blois Workshops.

- [253] FP420 Collaboration, M. Albrow et al., "FP420: A proposal to investigate the feasibility of installing proton tagging detectors in the 420 m region of the LHC," *CERN/LHCC 2005-025* (2005).
- [254] CDF Collaboration, A. A. Affolder et al., "Diffractive dijets with a leading antiproton in anti-p p collisions at $\sqrt{s} = 1800$ GeV," *Phys. Rev. Lett.* **84** (2000) 5043–5048.
- [255] A. De Roeck, V. A. Khoze, A. D. Martin, R. Orava, and M. G. Ryskin, "Ways to detect a light Higgs boson at the LHC," *Eur. Phys. J.* **C25** (2002) 391–403, arXiv:hep-ph/0207042.
- [256] V. Khoze, A. Martin, and M. Ryskin, "Prospects for new physics observations in diffractive processes at the LHC and Tevatron," *Eur. Phys. J.* **C23** (2002) 311–327, arXiv:hep-ph/0111078.
- [257] A. Kisselev, V. Petrov, and R. Ryutin, "5-dimensional quantum gravity effects in exclusive double diffractive events," *Phys. Lett.* **B630** (2005) 100–107. doi:10.1016/j.physletb.2005.09.059.
- [258] J. Monk and A. Pilkington, "ExHuME: A Monte Carlo event generator for exclusive diffraction," arXiv:hep-ph/0502077.
- [259] M. Boonekamp et al., "Monte-Carlo generators for central exclusive diffraction," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4montecarlo.pdf>.
- [260] R. Ryutin, "EDDE Monte Carlo event generator," arXiv:hep-ph/0409180.
- [261] A. Kaidalov, V. Khoze, A. Martin, and M. Ryskin, "Central exclusive diffractive production as a spin parity analyser: From hadrons to Higgs," *Eur. Phys. J.* **C31** (2003) 387–396, arXiv:hep-ph/0307064.
- [262] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, "MSSM Higgs boson searches at the Tevatron and the LHC: Impact of different benchmark scenarios," *Eur. Phys. J.* **C45** (2006) 797–814, arXiv:hep-ph/0511023.
- [263] J. R. Ellis, J. S. Lee, and A. Pilaftsis, "Diffraction as a CP and lineshape analyzer for MSSM Higgs bosons at the LHC," *Phys. Rev.* **D71** (2005) 075007, arXiv:hep-ph/0502251.
- [264] V. Petrov, R. Ryutin, A. Sobol, and J.-P. Guillaud, "Azimuthal angular distributions in EDDE as spin-parity analyser and glueball filter for LHC," *JHEP* **06** (2005) 007, arXiv:hep-ph/0409118.
- [265] K. Piotrkowski, "Tagging two-photon production at the LHC," *Phys. Rev.* **D63** (2001) 071502, arXiv:hep-ex/0009065.
- [266] V. Budnev, I. Ginzburg, G. Meledin, and V. Serbo, "The Two photon particle production mechanism. Physical problems. Applications. Equivalent photon approximation," *Phys. Rept.* **15** (1974) 181–281. doi:10.1016/0370-1573(75)90009-5.

- [267] V. Khoze, A. Martin, and M. Ryskin, "Photon-exchange processes at hadron colliders as a probe of the dynamics of diffraction," *Eur. Phys. J.* **C24** (2002) 459–468, arXiv:hep-ph/0201301. doi:10.1007/s10052-002-0964-4.
- [268] A. R. White, "The physics of a sextet quark sector," *Phys. Rev.* **D72** (2005) 036007, arXiv:hep-ph/0412062.
- [269] K. Piotrkowski, "High energy two-photon interactions at the LHC," arXiv:hep-ex/0201027.
- [270] N. N. Kalmykov, S. S. Ostapchenko, and A. I. Pavlov, "Quark-gluon string model and EAS simulation problems at ultra-high energies," *Nucl. Phys. Proc. Suppl.* **52B** (1997) 17–28.
- [271] R. Engel, T. K. Gaisser, T. Stanev, and P. Lipari, "Air shower calculations with the new version of SIBYLL." Prepared for 26th International Cosmic Ray Conference (ICRC 99), Salt Lake City, Utah, 17-25 Aug 1999.
- [272] S. Roesler, R. Engel, and J. Ranft, "The Event generator DPMJET-III at cosmic ray energies." Prepared for 27th International Cosmic Ray Conference (ICRC 2001), Hamburg, Germany, 7-15 Aug 2001.
- [273] F. Karsch, "Lattice QCD at high temperature and density," *Lect. Notes Phys.* **583** (2002) 209–249, arXiv:hep-lat/0106019.
- [274] E. Iancu and R. Venugopalan, "The color glass condensate and high energy scattering in QCD," arXiv:hep-ph/0303204.
- [275] D. J. Schwarz, "The first second of the universe," *Annalen Phys.* **12** (2003) 220–270, arXiv:astro-ph/0303574.
- [276] A. O. Starinets, "Transport coefficients of strongly coupled gauge theories: Insights from string theory," arXiv:nucl-th/0511073.
- [277] R. Bonciani, S. Catani, M. L. Mangano, and P. Nason, "NLL resummation of the heavy-quark hadroproduction cross-section," *Nucl. Phys.* **B529** (1998) 424–450, arXiv:hep-ph/9801375.
- [278] M. Davids et al., "Measurement of top-pair cross section and top-quark mass in the di-lepton and full-hadronic channels with CMS," *CMS Note* **2006/077** (2006).
- [279] S. Gennai et al., "Tau jet reconstruction and tagging at High Level Trigger and off-line," *CMS Note* **2006/028** (2006).
- [280] J. D'Hondt, J. Heyninck, and S. Lowette, "Measurement of the cross section of single leptonic $t\bar{t}$ events," *CMS Note* **2006/064** (2006).
- [281] C. Buttar et al., "Les Houches Physics at TeV Colliders 2005, Standard Model and Higgs working group: Summary report," arXiv:hep-ph/0604120.
- [282] V. Konopliyanikov, O. Kodolova, and A. Ulyanov, "Jet Calibration using γ +jet Events in the CMS Detector," *CMS Note* **2006/042** (2006).
- [283] J. D'Hondt et al., "Electron and muon reconstruction in single leptonic $t\bar{t}$ events," *CMS Note* **2006/024** (2006).

- [284] CMS Collaboration, D. Acosta et al., "CMS Coll. Physics TDR Vol.I, Section 12.2.8," *CERN/LHCC 2006-001* (2006).
- [285] S. Lowette, J. D'Hondt, J. Heyninck, and P. Vanlaer, "Offline Calibration of b-jet Identification Efficiency," CERN-CMS-NOTE-2006-013.
- [286] J. D'Hondt, S. Lowette, J. Heyninck, and S. Kassermann, "Light quark jet energy scale calibration using the W mass constraint in single-leptonic $t\bar{t}$ events," CERN-CMS-NOTE-2006-025.
- [287] The CDF Collaboration Collaboration, D. Acosta et al., "Measurement of the $t\bar{t}$ cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV using kinematic characteristics of lepton plus jets events," *Phys. Rev. D* **71** (2005) 072005. doi:10.1103/PhysRevD.71.072005.
- [288] DØ Collaboration, V. Abazov et al., "Measurement of the $t\bar{t}$ cross section in $p\bar{p}$ collisions at $\sqrt{s}=1.96$ TeV using kinematic characteristics of lepton plus jets events," *Phys. Lett. B* **626** (2005) 55. doi:10.1016/j.physletb.2005.08.105.
- [289] M. Vos and F. Palla, "B-tagging in the High Level Trigger," *CMS Note 2006-030* (2006).
- [290] CDF and DØ Collaboration, D. Wicke, "Top pair production cross-section measurement in the all-hadronic channel at CDF and DØ," *Int. J. Mod. Phys. A* **20** (2005) 3183–3186, arXiv:hep-ex/0411009. doi:10.1142/S0217751X05026091.
- [291] J. D'Hondt, J. Heyninck, and S. Lowette, "Top Quark mass measurement in single-leptonic $t\bar{t}$ events," *CMS Note 2006/066* (2006).
- [292] CDF Collaboration, F. Abe et al., "First Observation of the All-Hadronic Decay of $t\bar{t}$ pairs," *Phys. Rev. Lett.* **79** (1997) 1992–1997. doi:10.1103/PhysRevLett.79.1992.
- [293] Tevatron Electroweak Working Group Collaboration, "Combination of CDF and DØ results on the mass of the top quark," arXiv:hep-ex/0603039.
- [294] A. Kharchilava, "Top mass determination in leptonic final states with J/ψ ," *Phys. Lett. B* **476** (2000) 73–78, arXiv:hep-ph/9912320.
- [295] P. Grenier, "," *ATLAS Physics Note 2001-023* (2001).
- [296] R. Chierici and A. Dierlamm, "Determination of the top mass in exclusive J/ψ decays," *CMS Note 2006-058* (2006).
- [297] Tevatron Electroweak Working Group Collaboration, "Combination of CDF and DØ results on the mass of the top quark," arXiv:hep-ex/0603039.
- [298] C. S. Hill, J. R. Incandela, and J. M. Lamb, "A method for measurement of the top quark mass using the mean decay length of b hadrons in t anti-t events," *Phys. Rev. D* **71** (2005) 054029, arXiv:hep-ex/0501043.
- [299] I. Borjanovic et al., "Investigation of top mass measurements with the ATLAS detector at LHC," *Eur. Phys. J. C* **39S2** (2005) 63–90, arXiv:hep-ex/0403021.
- [300] L. Lyons, D. Gibaut, and P. Clifford, "How to combine correlated estimates of a single physical quantity," *Nucl. Instrum. Meth. A* **270** (1988) 110.

- [301] A. Valassi, "Combining correlated measurements of several different physical quantities," *Nucl. Instrum. Meth.* **A500** (2003) 391–405.
- [302] G. Mahlon and S. Parke, "Angular Correlation in Top Quark Pair Production and Decay at Hadron Collider," *Phys. Rev.* **D 53** (1996) 4886–4896, arXiv:hep-ph/9512264.
- [303] T. Stelzer and S. Willenbrock, "Spin correlation in top-quark production at hadron collider," *Phys. Lett.* **B 374** (1996) 169–172, arXiv:hep-ph/9512292.
- [304] A. Brandenburg, "Spin-Spin Correlations of Top Quark Pairs at Hadron Colliders," *Phys. Lett.* **B 387** (1996) 626–632, arXiv:hep-ph/9606379.
- [305] M. Baarmand, H. Mermerkaya, and I. Vodopiyanov, "Measurement of Spin Correlation in Top Quark Pair Production in Semi-Leptonic Final State," *CMS Note* **2006/111** (2006).
- [306] T. Tait, "The tW^- mode of single top production," *Phys. Rev.* **D61** (2000) 034001, arXiv:hep-ph/9909352. doi:10.1103/PhysRevD.61.034001.
- [307] A. Belyaev and E. Boos, "Single top quark $tW + X$ production at the LHC: A closer look," *Phys. Rev.* **D 63** (2001) 034012, arXiv:hep-ph/0003260.
- [308] S. Zhu, "Next-to-leading order QCD corrections to $b g \rightarrow t W^-$ at the CERN Large Hadron Collider," *Phys. Lett.* **B 524** (2002) 283–288.
- [309] E. Boos, V. Bunichev, L. Dudko, V. Savrin, and A. Sherstnev, "A simulation method of the Electroweak Top Quark Production Events in the NLO Approximation. A Monte-Carlo Generator 'SingleTop'," 2005. Abstract available at http://www.npi.msu.su/eng/science.php3?sec=preprint&ref_pp=1262.
- [310] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 9.1.2: Global muon reconstruction," *CERN/LHCC* **2006-001** (2006) 333.
- [311] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 9.3: Muon identification," *CERN/LHCC* **2006-001** (2006) 351.
- [312] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 10.4: Electron reconstruction and selection," *CERN/LHCC* **2006-001** (2006) 390.
- [313] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 11.2.1: Iterative cone," *CERN/LHCC* **2006-001** (2006) 408.
- [314] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 11.6.3: γ +jet events," *CERN/LHCC* **2006-001** (2006) 423.
- [315] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 12.2: b-tagging tools," *CERN/LHCC* **2006-001** (2006) 461.
- [316] V. Abramov et al., "Selection of single top events with the CMS detector at LHC," *CMS Note* **2006/084** (2006).
- [317] P. Yeh et al., "Search for W-associated Production of Single Top Quarks in CMS," *CMS Note* **2006/086** (2006).

- [318] J. Campbell, R. K. Ellis, and D. L. Rainwater, "Next-to-leading order QCD predictions for $W + 2\text{jet}$ and $Z + 2\text{jet}$ production at the CERN LHC," *Phys. Rev.* **D68** (2003) 094021. doi:10.1103/PhysRevD.68.094021.
- [319] CMS Collaboration, D. Acosta et al., "CMS Coll. Physics TDR Vol.I, Section 11.6.5," *CERN/LHCC 2006-001* (2006).
- [320] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 8.5: Sources of systematic effects," *CERN/LHCC 2006-001* (2006) 229.
- [321] R. A. Fisher, "The use of multiple measurements in taxonomic problems," *Annals of Eugenics* **7** (1936) 179–188.
- [322] B. Mele, S. Petrarca, and A. Soddu, "A new evaluation of the $t \rightarrow cH$ decay width in the standard model," *Phys. Lett.* **B435** (1998) 401–406, arXiv:hep-ph/9805498.
- [323] C.-S. Huang, X.-H. Wu, and S.-H. Zhu, "Top-charm associated production at high energy $e^+ e^-$ colliders in standard model," *Phys. Lett.* **B452** (1999) 143–149, arXiv:hep-ph/9901369.
- [324] G. M. de Divitiis, R. Petronzio, and L. Silvestrini, "Flavour changing top decays in supersymmetric extensions of the standard model," *Nucl. Phys.* **B504** (1997) 45–60, arXiv:hep-ph/9704244.
- [325] J. Guasch and J. Sola, "FCNC top quark decays: A door to SUSY physics in high luminosity colliders?," *Nucl. Phys.* **B562** (1999) 3–28, arXiv:hep-ph/9906268.
- [326] G. Eilam, A. Gemintern, T. Han, J. M. Yang, and X. Zhang, "Top quark rare decay $t \rightarrow ch$ in R-parity-violating SUSY," *Phys. Lett.* **B510** (2001) 227–235, arXiv:hep-ph/0102037. doi:10.1016/S0370-2693(01)00598-6.
- [327] C. S. Li, X.-m. Zhang, and S. H. Zhu, "SUSY-QCD effect on top charm associated production at linear collider," *Phys. Rev.* **D60** (1999) 077702, arXiv:hep-ph/9904273.
- [328] S. Bejar, J. Guasch, and J. Sola, "FCNC top quark decays beyond the standard model," arXiv:hep-ph/0101294.
- [329] R. A. Diaz, R. Martinez, and J. Alexis Rodriguez, "The rare decay $t \rightarrow c\gamma$ in the general 2HDM type III," arXiv:hep-ph/0103307.
- [330] T. Han and J. L. Hewett, "Top charm associated production in high energy $e^+ e^-$ collisions," *Phys. Rev.* **D60** (1999) 074015, arXiv:hep-ph/9811237.
- [331] F. del Aguila, J. A. Aguilar-Saavedra, and R. Miquel, "Constraints on top couplings in models with exotic quarks," *Phys. Rev. Lett.* **82** (1999) 1628–1631, arXiv:hep-ph/9808400.
- [332] J. A. Aguilar-Saavedra and B. M. Nobre, "Rare top decays $t \rightarrow c\gamma$, $t \rightarrow cg$ and CKM unitarity," *Phys. Lett.* **B553** (2003) 251–260, arXiv:hep-ph/0210360.
- [333] CDF and DØ Collaboration, M. Paulini, "Heavy flavor physics from top to bottom," arXiv:hep-ex/9701019.

- [334] CDF Collaboration, F. Abe et al., "Search for Flavor-Changing Neutral Current Decays of the Top Quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **80** (1998) 2525–2530. doi:10.1103/PhysRevLett.80.2525.
- [335] A. Heinson, "Future top physics at the Tevatron and LHC," arXiv:hep-ex/9605010.
- [336] J. A. Aguilar-Saavedra and G. C. Branco, "Probing top flavour-changing neutral scalar couplings at the CERN LHC," *Phys. Lett.* **B495** (2000) 347–356, arXiv:hep-ph/0004190.
- [337] J. A. Aguilar-Saavedra, "Top flavour-changing neutral coupling signals at a linear collider," *Phys. Lett.* **B502** (2001) 115–124, arXiv:hep-ph/0012305.
- [338] J. A. Aguilar-Saavedra and T. Riemann, "Probing top flavor-changing neutral couplings at TESLA," arXiv:hep-ph/0102197.
- [339] K. Karafasoulis et al., "Study of Flavour Changing Neutral Currents in Top Quark Decays with the CMS Detector," *CMS Note* **2006/093** (2006).
- [340] G. Dissertori et al., "How accurately can we count the number of $pp \rightarrow ZX$ and $pp \rightarrow WX$ events using decays to electrons," *CMS Note* **2006/124** (2006).
- [341] J. Alcaraz, "Measurement of $Z \rightarrow \mu^+ \mu^-$ and $W \rightarrow \mu\nu$ rates in CMS," *CMS Note* **2006/082** (2006).
- [342] S. Frixione and B. R. Webber, "The MC@NLO 3.2 event generator," arXiv:hep-ph/0601192.
- [343] For further details see Proceedings of the 2004/2005 HERA-LHC workshop.
- [344] M. Dittmar, F. Pauss, and D. Zuercher, "Towards a precise parton luminosity determination at the CERN LHC," *Phys. Rev.* **D56** (1997) 7284–7290, arXiv:hep-ex/9705004.
- [345] S. D. Drell and T.-M. Yan, "Massive Lepton Pair Production in Hadron-Hadron Collisions at High-Energies," *Phys. Rev. Lett.* **25** (1970) 316–320. doi:10.1103/PhysRevLett.25.316.
- [346] I. Belotelov et al., "Study of Drell-Yan Di-muon Production with the CMS Detector," *CMS Note* **2006/123** (2006).
- [347] R. Hamberg, W. L. van Neerven, and T. Matsuura, "A Complete calculation of the order α_s^2 correction to the Drell-Yan K -factor," *Nucl. Phys.* **B359** (1991) 343–405. doi:10.1016/0550-3213(91)90064-5.
- [348] D. Bourilkov, "Compositeness Search with Di-muons in CMS," *CMS Note* **2006/085** (2006).
- [349] U. Baur and D. Wackerth, "Electroweak radiative corrections to weak boson production at hadron colliders," *Nucl. Phys. Proc. Suppl.* **116** (2003) 159–163, arXiv:hep-ph/0211089.

- [350] D. Bourilkov, "Study of Parton Density Function Uncertainties with LHAPDF and PYTHIA at LHC," arXiv:hep-ph/0305126. Prepared for the LHC/LC Study Group Meeting, Geneva, Switzerland, 9 May 2003.
- [351] V. Buege et al., "Prospects for the precision measurement of the W mass with the CMS detector," *CMS Note* **2006-061** (2006).
- [352] W. T. Giele and S. Keller, "Determination of W boson properties at hadron colliders," *Phys. Rev.* **D57** (1998) 4433–4440, arXiv:hep-ph/9704419.
- [353] V. Brigljević et al., "Study of Di-Boson Production with the CMS Detector at the LHC," *CMS Note* **2006/108** (2006).
- [354] A. Pukhov et al., "CompHEP: A package for evaluation of Feynman diagrams and integration over multi-particle phase space. User's manual for version 33," arXiv:hep-ph/9908288.
- [355] P. W. Higgs, "Broken symmetries, massless particles and gauge fields," *Phys. Lett.* **12** (1964) 132–133.
- [356] P. W. Higgs, "Spontaneous Symmetry Breakdown Without Massless Bosons," *Phys. Rev.* **145** (1966) 1156–1163.
- [357] F. Englert and R. Brout, "Broken Symmetry and the Mass of Gauge Vector Mesons," *Phys. Rev. Lett.* **13** (1964) 321–322.
- [358] G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, "Global Conservation Laws and Massless Particles," *Phys. Rev. Lett.* **13** (1964) 585–587.
- [359] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, "Uniqueness of spontaneously broken gauge theories," *Phys. Rev. Lett.* **30** (1973) 1268–1270.
doi:10.1103/PhysRevLett.30.1268.
- [360] J. M. Cornwall, D. N. Levin, and G. Tiktopoulos, "Derivation of Gauge Invariance from High-Energy Unitarity Bounds on the S - Matrix," *Phys. Rev.* **D10** (1974) 1145.
- [361] C. H. Llewellyn Smith, "High-Energy Behavior and Gauge Symmetry," *Phys. Lett.* **B46** (1973) 233–236.
- [362] S. D. Joglekar, "S-matrix derivation of the Weinberg model," *Ann. Phys.* **83** (1974) 427.
- [363] M. Veltman, "Perturbation theory of massive Yang-Mills fields," *Nucl. Phys.* **B7** (1968) 637–650.
- [364] G. 't Hooft, "Renormalization of Massless Yang-Mills Fields," *Nucl. Phys.* **B33** (1971) 173–199.
- [365] G. 't Hooft, "Renormalizable Lagrangians for Massive Yang-Mills Fields," *Nucl. Phys.* **B35** (1971) 167–188.
- [366] G. 't Hooft and M. J. G. Veltman, "Regularization and Renormalization of Gauge Fields," *Nucl. Phys.* **B44** (1972) 189–213.
- [367] G. 't Hooft and M. J. G. Veltman, "Combinatorics of gauge fields," *Nucl. Phys.* **B50** (1972) 318–353.

- [368] A. Djouadi, "The anatomy of electro-weak symmetry breaking. I: The Higgs boson in the standard model," *arXiv:hep-ph/0503172*.
- [369] A. Djouadi, "The anatomy of electro-weak symmetry breaking. II: The Higgs bosons in the minimal supersymmetric model," *arXiv:hep-ph/0503173*.
- [370] N. Cabibbo, L. Maiani, G. Parisi, and R. Petronzio, "Bounds on the Fermions and Higgs Boson Masses in Grand Unified Theories," *Nucl. Phys.* **B158** (1979) 295.
- [371] M. S. Chanowitz, M. A. Furman, and I. Hinchliffe, "Weak Interactions of Ultraheavy Fermions," *Phys. Lett.* **B78** (1978) 285.
- [372] R. A. Flores and M. Sher, "Upper Limits to Fermion Masses in The Glashow-Weinberg-Salam Model," *Phys. Rev.* **D27** (1983) 1679.
- [373] M. Lindner, "Implications of Triviality for the Standard Model," *Zeit. Phys.* **C31** (1986) 295.
- [374] M. Sher, "Electroweak Higgs Potentials and Vacuum Stability," *Phys. Rept.* **179** (1989) 273–418.
- [375] M. Sher, "Precise vacuum stability bound in the standard model," *Phys. Lett.* **B317** (1993) 159–163, *arXiv:hep-ph/9307342*.
- [376] G. Altarelli and G. Isidori, "Lower limit on the Higgs mass in the standard model: An Update," *Phys. Lett.* **B337** (1994) 141–144.
- [377] J. Espinosa and M. Quiros, "Improved metastability bounds on the standard model Higgs mass," *Phys. Lett.* **B353** (1995) 257–266, *arXiv:hep-ph/9504241*.
- [378] A. Hasenfratz, K. Jansen, C. B. Lang, T. Neuhaus, and H. Yoneyama, "The Triviality Bound of the Four Component Φ^4 Model," *Phys. Lett.* **B199** (1987) 531.
- [379] J. Kuti, L. Lin, and Y. Shen, "Upper Bound on the Higgs Mass in the Standard Model," *Phys. Rev. Lett.* **61** (1988) 678.
- [380] M. Luscher and P. Weisz, "Scaling Laws and Triviality Bounds in the Lattice ϕ^4 Theory (III). n -Component Model," *Nucl. Phys.* **B318** (1989) 705. doi:10.1016/0550-3213(89)90637-8.
- [381] **Higgs Working Group** Collaboration, M. Carena et al., "Report of the Tevatron Higgs working group," *arXiv:hep-ph/0010338*.
- [382] ATLAS Collaboration, "Technical Design Report," *CERN/LHCC 94-14* (1994). ATLAS TDR 14.
- [383] S. Abdullin et al., "Summary of the CMS potential for the Higgs boson discovery," *Eur. Phys. J.* **C39S2** (2005) 41–61. doi:10.1140/epjcd/s2004-02-003-9.
- [384] E. Braaten and J. P. Leveille, "Higgs Boson Decay and the Running Mass," *Phys. Rev.* **D22** (1980) 715.
- [385] N. Sakai, "Perturbative QCD Corrections to the Hadronic Decay Width of the Higgs Boson," *Phys. Rev.* **D22** (1980) 2220.

- [386] T. Inami and T. Kubota, "Renormalization Group Estimate of the Hadronic Decay Width of the Higgs Boson," *Nucl. Phys.* **B179** (1981) 171.
- [387] S. G. Gorishnii, A. L. Kataev, and S. A. Larin, "The Width of Higgs Boson Decay into Hadrons: Three Loop Corrections of Strong Interactions," *Sov. J. Nucl. Phys.* **40** (1984) 329–334.
- [388] M. Drees and K.-i. Hikasa, "Heavy Quark Thresholds in Higgs Physics," *Phys. Rev.* **D41** (1990) 1547.
- [389] M. Drees and K.-i. Hikasa, "Note on QCD Corrections to Hadronic Higgs Decay," *Phys. Lett.* **B240** (1990) 455.
- [390] K. G. Chetyrkin, "Correlator of the quark scalar currents and $\Gamma(\text{tot})(H \rightarrow \text{hadrons})$ at $\mathcal{O}(\alpha_s^3)$ in pQCD," *Phys. Lett.* **B390** (1997) 309–317, arXiv:hep-ph/9608318.
- [391] J. Fleischer and F. Jegerlehner, "Radiative Corrections to Higgs Decays in the Extended Weinberg-Salam Model," *Phys. Rev.* **D23** (1981) 2001–2026.
- [392] D. Y. Bardin, B. M. Vilensky, and P. K. Khristova, "Calculation of the Higgs boson decay width into fermion pairs," *Sov. J. Nucl. Phys.* **53** (1991) 152–158.
- [393] A. Dabelstein and W. Hollik, "Electroweak corrections to the fermionic decay width of the standard Higgs boson," *Z. Phys.* **C53** (1992) 507–516.
- [394] B. A. Kniehl, "Radiative corrections for $H \rightarrow f\bar{f}(\gamma)$ in the standard model," *Nucl. Phys.* **B376** (1992) 3–28. doi:10.1016/0550-3213(92)90065-J.
- [395] H.-Q. Zheng and D.-D. Wu, "First order QCD corrections to the decay of the Higgs boson into two photons," *Phys. Rev.* **D42** (1990) 3760–3763.
- [396] A. Djouadi, M. Spira, J. van der Bij, and P. Zerwas, "QCD corrections to gamma gamma decays of Higgs particles in the intermediate mass range," *Phys. Lett.* **B257** (1991) 187–190.
- [397] S. Dawson and R. P. Kauffman, "QCD corrections to $H \rightarrow \gamma\gamma$," *Phys. Rev.* **D47** (1993) 1264–1267.
- [398] A. Djouadi, M. Spira, and P. Zerwas, "Two photon decay widths of Higgs particles," *Phys. Lett.* **B311** (1993) 255–260, arXiv:hep-ph/9305335.
- [399] K. Melnikov and O. I. Yakovlev, "Higgs \rightarrow two photon decay: QCD radiative correction," *Phys. Lett.* **B312** (1993) 179–183, arXiv:hep-ph/9302281.
- [400] M. Inoue, R. Najima, T. Oka, and J. Saito, "QCD corrections to two photon decay of the Higgs boson and its reverse process," *Mod. Phys. Lett.* **A9** (1994) 1189–1194.
- [401] M. Steinhauser, "Corrections of $\mathcal{O}(\alpha_s^2)$ to the decay of an intermediate-mass Higgs boson into two photons," arXiv:hep-ph/9612395.
- [402] A. Djouadi, P. Gambino, and B. A. Kniehl, "Two-loop electroweak heavy-fermion corrections to Higgs-boson production and decay," *Nucl. Phys.* **B523** (1998) 17–39, arXiv:hep-ph/9712330.

- [403] U. Aglietti, R. Bonciani, G. Degrossi, and A. Vicini, "Two-loop light fermion contribution to Higgs production and decays," *Phys. Lett.* **B595** (2004) 432–441, arXiv:hep-ph/0404071.
- [404] G. Degrossi and F. Maltoni, "Two-loop electroweak corrections to the Higgs-boson decay $H \rightarrow \gamma \gamma$," arXiv:hep-ph/0504137.
- [405] B. A. Kniehl, "Radiative corrections for $H \rightarrow ZZ$ in the standard model," *Nucl. Phys.* **B352** (1991) 1–26. doi:10.1016/0550-3213(91)90126-I.
- [406] B. A. Kniehl, "Radiative corrections for $H \rightarrow W^+W^-(\gamma)$ in the standard model," *Nucl. Phys.* **B357** (1991) 439–466. doi:10.1016/0550-3213(91)90476-E.
- [407] H. Georgi, S. Glashow, M. Machacek, and D. Nanopoulos, "Higgs Bosons from Two Gluon Annihilation in Proton Proton Collisions," *Phys. Rev. Lett.* **40** (1978) 692. doi:10.1103/PhysRevLett.40.692.
- [408] D. Graudenz, M. Spira, and P. M. Zerwas, "QCD corrections to Higgs boson production at proton proton colliders," *Phys. Rev. Lett.* **70** (1993) 1372–1375. doi:10.1103/PhysRevLett.70.1372.
- [409] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, "SUSY Higgs production at proton colliders," *Phys. Lett.* **B318** (1993) 347–353. doi:10.1016/0370-2693(93)90138-8.
- [410] M. Spira, A. Djouadi, D. Graudenz, and P. Zerwas, "Higgs boson production at the LHC," *Nucl. Phys.* **B453** (1995) 17–82, arXiv:hep-ph/9504378. doi:10.1016/0550-3213(95)00379-7.
- [411] M. Kramer, E. Laenen, and M. Spira, "Soft gluon radiation in Higgs boson production at the LHC," *Nucl. Phys.* **B511** (1998) 523–549, arXiv:hep-ph/9611272. doi:10.1016/S0550-3213(97)00679-2.
- [412] A. Djouadi, M. Spira, and P. Zerwas, "Production of Higgs bosons in proton colliders: QCD corrections," *Phys. Lett.* **B264** (1991) 440–446.
- [413] S. Dawson, "Radiative corrections to Higgs boson production," *Nucl. Phys.* **B359** (1991) 283–300.
- [414] R. P. Kauffman and W. Schaffer, "QCD corrections to production of Higgs pseudoscalars," *Phys. Rev.* **D49** (1994) 551–554, arXiv:hep-ph/9305279.
- [415] S. Dawson and R. Kauffman, "QCD corrections to Higgs boson production: nonleading terms in the heavy quark limit," *Phys. Rev.* **D49** (1994) 2298–2309, arXiv:hep-ph/9310281.
- [416] R. V. Harlander and W. B. Kilgore, "Next-to-next-to-leading order Higgs production at hadron colliders," *Phys. Rev. Lett.* **88** (2002) 201801, arXiv:hep-ph/0201206.
- [417] R. V. Harlander and W. B. Kilgore, "Production of a pseudo-scalar Higgs boson at hadron colliders at next-to-next-to leading order," *JHEP* **10** (2002) 017, arXiv:hep-ph/0208096.

- [418] C. Anastasiou and K. Melnikov, "Higgs boson production at hadron colliders in NNLO QCD," *Nucl. Phys.* **B646** (2002) 220–256, arXiv:hep-ph/0207004.
- [419] V. Ravindran, J. Smith, and W. L. van Neerven, "NNLO corrections to the total cross section for Higgs boson production in hadron hadron collisions," *Nucl. Phys.* **B665** (2003) 325–366, arXiv:hep-ph/0302135.
- [420] S. Catani, D. de Florian, M. Grazzini, and P. Nason, "Soft-gluon resummation for Higgs boson production at hadron colliders," *JHEP* **07** (2003) 028, arXiv:hep-ph/0306211.
- [421] A. Djouadi and P. Gambino, "Leading electroweak correction to Higgs boson production at proton colliders," *Phys. Rev. Lett.* **73** (1994) 2528–2531, arXiv:hep-ph/9406432.
- [422] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, "Virtual top-quark effects on the $H \rightarrow b$ anti- b decay at next-to-leading order in QCD," *Phys. Rev. Lett.* **78** (1997) 594–597, arXiv:hep-ph/9610456.
- [423] K. G. Chetyrkin, B. A. Kniehl, and M. Steinhauser, "Three-loop $O(\alpha(s)^2 G(F) M(t)^2)$ corrections to hadronic Higgs decays," *Nucl. Phys.* **B490** (1997) 19–39, arXiv:hep-ph/9701277.
- [424] A. Ghinculov and J. J. van der Bij, "The Higgs resonance shape in gluon fusion: Heavy Higgs effects," *Nucl. Phys.* **B482** (1996) 59–72, arXiv:hep-ph/9511414.
- [425] R. K. Ellis, I. Hinchliffe, M. Soldate, and J. J. van der Bij, "Higgs Decay to $\tau^+ \tau^-$: A Possible Signature of Intermediate Mass Higgs Bosons at the SSC," *Nucl. Phys.* **B297** (1988) 221.
- [426] U. Baur and E. W. N. Glover, "Higgs Boson Production at Large Transverse Momentum in Hadronic Collisions," *Nucl. Phys.* **B339** (1990) 38–66.
- [427] C. R. Schmidt, " $H \rightarrow ggg(gq\bar{q})$ at two loops in the large- M_t limit," *Phys. Lett.* **B413** (1997) 391–395. doi:10.1016/S0370-2693(97)01102-7.
- [428] D. de Florian, M. Grazzini, and Z. Kunszt, "Higgs production with large transverse momentum in hadronic collisions at next-to-leading order," *Phys. Rev. Lett.* **82** (1999) 5209–5212, arXiv:hep-ph/9902483.
- [429] V. Ravindran, J. Smith, and W. L. Van Neerven, "Next-to-leading order QCD corrections to differential distributions of Higgs boson production in hadron hadron collisions," *Nucl. Phys.* **B634** (2002) 247–290, arXiv:hep-ph/0201114.
- [430] C. J. Glosser and C. R. Schmidt, "Next-to-leading corrections to the Higgs boson transverse momentum spectrum in gluon fusion," *JHEP* **12** (2002) 016, arXiv:hep-ph/0209248.
- [431] C. Anastasiou, K. Melnikov, and F. Petriello, "Fully differential Higgs boson production and the di-photon signal through next-to-next-to-leading order," *Nucl. Phys.* **B724** (2005) 197–246, arXiv:hep-ph/0501130.
- [432] S. Catani, E. D’Emilio, and L. Trentadue, "The Gluon Form-Factor to Higher Orders: Gluon Gluon Annihilation at Small Q-Transverse," *Phys. Lett.* **B211** (1988) 335–342.

- [433] I. Hinchliffe and S. F. Novaes, "On the Mean Transverse Momentum of Higgs Bosons at the SSC," *Phys. Rev.* **D38** (1988) 3475–3480.
- [434] R. P. Kauffman, "Higgs boson $p(T)$ in gluon fusion," *Phys. Rev.* **D44** (1991) 1415–1425.
- [435] R. Kauffman, "Higher order corrections to Higgs boson $p(T)$," *Phys. Rev.* **D45** (1992) 1512–1517.
- [436] C. Balazs and C. P. Yuan, "Higgs boson production at the LHC with soft gluon effects," *Phys. Lett.* **B478** (2000) 192–198, arXiv:hep-ph/0001103.
- [437] E. L. Berger and J.-w. Qiu, "Differential cross section for Higgs boson production including all-orders soft gluon resummation," *Phys. Rev.* **D67** (2003) 034026, arXiv:hep-ph/0210135.
- [438] A. Kulesza and W. J. Stirling, "Non-perturbative effects and the resummed Higgs transverse momentum distribution at the LHC," *JHEP* **12** (2003) 056, arXiv:hep-ph/0307208.
- [439] A. Kulesza, G. Sterman, and W. Vogelsang, "Joint resummation for Higgs production," *Phys. Rev.* **D69** (2004) 014012, arXiv:hep-ph/0309264.
- [440] A. Gawron and J. Kwiecinski, "Resummation effects in Higgs boson transverse momentum distribution within the framework of unintegrated parton distributions," *Phys. Rev.* **D70** (2004) 014003, arXiv:hep-ph/0309303.
- [441] G. Watt, A. Martin, and M. Ryskin, "Unintegrated parton distributions and electroweak boson production at hadron colliders," *Phys. Rev.* **D70** (2004) 014012, arXiv:hep-ph/0309096.
- [442] A. Lipatov and N. Zotov, "Higgs boson production at hadron colliders in the $k(T)$ -factorization approach," *Eur. Phys. J.* **C44** (2005) 559–566, arXiv:hep-ph/0501172.
- [443] D. de Florian and M. Grazzini, "Next-to-next-to-leading logarithmic corrections at small transverse momentum in hadronic collisions," *Phys. Rev. Lett.* **85** (2000) 4678–4681, arXiv:hep-ph/0008152.
- [444] S. Catani, D. De Florian, and M. Grazzini, "Higgs production at hadron colliders in (almost) NNLO QCD," in *Bologna 2001, Deep inelastic scattering*, pp. 518–521. Bologna, Italy, April, 2001. arXiv:hep-ph/0106049.
- [445] S. Catani, D. de Florian, and M. Grazzini, "Universality of non-leading logarithmic contributions in transverse momentum distributions," *Nucl. Phys.* **B596** (2001) 299–312, arXiv:hep-ph/0008184.
- [446] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, "The $q(T)$ spectrum of the Higgs boson at the LHC in QCD perturbation theory," *Phys. Lett.* **B564** (2003) 65–72, arXiv:hep-ph/0302104.
- [447] G. Bozzi, S. Catani, D. de Florian, and M. Grazzini, "Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC," *Nucl. Phys.* **B737** (2006) 73–120, arXiv:hep-ph/0508068.

- [448] R. N. Cahn and S. Dawson, "Production of Very Massive Higgs Bosons," *Phys. Lett.* **B136** (1984) 196.
- [449] K.-i. Hikasa, "Heavy Higgs Production in $e^+ e^-$ and $e^- e^-$ Collisions," *Phys. Lett.* **B164** (1985) 385.
- [450] G. Altarelli, B. Mele, and F. Pitolli, "Heavy Higgs Production at Future Colliders," *Nucl. Phys.* **B287** (1987) 205–224.
- [451] T. Han, G. Valencia, and S. Willenbrock, "Structure function approach to vector boson scattering in $p p$ collisions," *Phys. Rev. Lett.* **69** (1992) 3274–3277, [arXiv:hep-ph/9206246](https://arxiv.org/abs/hep-ph/9206246).
- [452] T. Figy, C. Oleari, and D. Zeppenfeld, "Next-to-leading order jet distributions for Higgs boson production via weak-boson fusion," *Phys. Rev.* **D68** (2003) 073005, [arXiv:hep-ph/0306109](https://arxiv.org/abs/hep-ph/0306109).
- [453] S. Glashow, D. Nanopoulos, and A. Yildiz, "Associated Production of Higgs Bosons and Z Particles," *Phys. Rev.* **D18** (1978) 1724–1727.
- [454] Z. Kunszt, Z. Trocsanyi, and W. J. Stirling, "Clear signal of intermediate mass Higgs boson production at LHC and SSC," *Phys. Lett.* **B271** (1991) 247–255.
- [455] T. Han and S. Willenbrock, "QCD correction to the $p p \rightarrow W H$ and $Z H$ total cross-sections," *Phys. Lett.* **B273** (1991) 167–172.
- [456] O. Brein, A. Djouadi, and R. Harlander, "NNLO QCD corrections to the Higgs-strahlung processes at hadron colliders," *Phys. Lett.* **B579** (2004) 149–156, [arXiv:hep-ph/0307206](https://arxiv.org/abs/hep-ph/0307206).
- [457] M. L. Ciccolini, S. Dittmaier, and M. Kramer, "Electroweak radiative corrections to associated $W H$ and $Z H$ production at hadron colliders," *Phys. Rev.* **D68** (2003) 073003, [arXiv:hep-ph/0306234](https://arxiv.org/abs/hep-ph/0306234).
- [458] R. Raitio and W. W. Wada, "Higgs Boson Production at Large Transverse Momentum in QCD," *Phys. Rev.* **D19** (1979) 941.
- [459] J. N. Ng and P. Zakarauskas, "A QCD Parton Calculation of Conjoined Production of Higgs Bosons and Heavy Flavors in p anti- p Collision," *Phys. Rev.* **D29** (1984) 876.
- [460] Z. Kunszt, "Associated Production of Heavy Higgs Boson with Top Quarks," *Nucl. Phys.* **B247** (1984) 339.
- [461] J. F. Gunion, "Associated top anti-top Higgs production as a large source of $W H$ events: Implications for Higgs detection in the lepton neutrino $\gamma \gamma$ final state," *Phys. Lett.* **B261** (1991) 510–517.
- [462] W. J. Marciano and F. E. Paige, "Associated production of Higgs bosons with t anti- t pairs," *Phys. Rev. Lett.* **66** (1991) 2433–2435.
- [463] W. Beenakker, S. Dittmaier, M. Kraemer, B. Pluemper, M. Spira, and P. Zerwas, "Higgs radiation off top quarks at the Tevatron and the LHC," *Phys. Rev. Lett.* **87** (2001) 201805, [arXiv:hep-ph/0107081](https://arxiv.org/abs/hep-ph/0107081).

- [464] S. Dawson, L. H. Orr, L. Reina, and D. Wackerth, "Associated top quark Higgs boson production at the LHC," *Phys. Rev.* **D67** (2003) 071503, arXiv:hep-ph/0211438.
- [465] S. Dawson and L. Reina, "QCD corrections to associated Higgs boson production," *Phys. Rev.* **D57** (1998) 5851–5859, arXiv:hep-ph/9712400.
- [466] D. Futyan, D. Fortin, and D. Giordano, "Search for the Standard Model Higgs Boson in the Two-Electron and Two-Muon Final State with CMS," *CMS Note* **2006/136** (2006).
- [467] S. Abdullin et al., "Search Strategy for the Standard Model Higgs Boson in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ Decay Channel using $M(4\mu)$ -Dependent Cuts," *CMS Note* **2006/122** (2006).
- [468] P. Bartalini et al., "NLO vs. LO: kinematical differences for signal and background in the $H \rightarrow ZZ^{(*)} \rightarrow 4\mu$ analysis," *CMS Note* (2006/130).
- [469] S. Baffioni et al., "Discovery potential for the SM Higgs boson in the $H \rightarrow ZZ^{(*)} \rightarrow e^+e^-e^+e^-$ decay channel," *CMS Note* **2006/115** (2006).
- [470] M. Dittmar and H. K. Dreiner, "How to find a Higgs boson with a mass between 155-GeV to 180-GeV at the LHC," *Phys. Rev.* **D55** (1997) 167–172, arXiv:hep-ph/9608317.
- [471] G. Davatz, M. Dittmar, and A.-S. Giolo-Nicollerat, "Standard Model Higgs Discovery Potential of CMS in $H \rightarrow WW \rightarrow \ell\nu\ell\nu$ Channel," *CMS Note* **2006/047** (2006).
- [472] G. Davatz, G. Dissertori, M. Dittmar, M. Grazzini, and F. Pauss, "Effective K-factors for $gg \rightarrow H \rightarrow WW \rightarrow \ell\nu\ell\nu$ at the LHC," *JHEP* **05** (2004) 009, arXiv:hep-ph/0402218.
- [473] J. Campbell and F. Tramontano, "Next-to-leading order corrections to Wt production and decay," *Nucl. Phys.* **B726** (2005) 109–130, arXiv:hep-ph/0506289.
- [474] CMS Collaboration, F. Beaudette et al., "Search for a Light Standard Model Higgs Boson in the $H \rightarrow WW^{(*)} \rightarrow e^+\nu e^-\bar{\nu}$ Channel," *CMS Note* **2006/114** (2006).
- [475] T. Plehn, D. L. Rainwater, and D. Zeppenfeld, "A method for identifying $H \rightarrow \tau\tau \rightarrow e^\pm\mu^\mp p_T$ at the CERN LHC," *Phys. Rev.* **D61** (2000) 093005. doi:10.1103/PhysRevD.61.093005.
- [476] D. L. Rainwater, D. Zeppenfeld, and K. Hagiwara, "Searching for $H \rightarrow \tau\tau$ in weak boson fusion at the LHC," *Phys. Rev.* **D59** (1999) 014037. doi:10.1103/PhysRevD.59.014037.
- [477] D. Cavalli et al., "The Higgs working group: Summary report," arXiv:hep-ph/0203056.
- [478] D. Zeppenfeld, R. Kinnunen, A. Nikitenko, and E. Richter-Was, "Measuring Higgs boson couplings at the LHC," *Phys. Rev.* **D62** (2000) 013009, arXiv:hep-ph/0002036.
- [479] M. Dührssen et al., "Extracting Higgs boson couplings from LHC data," *Phys. Rev.* **D70** (2004) 113009, arXiv:hep-ph/0406323.

- [480] T. Plehn, D. L. Rainwater, and D. Zeppenfeld, "Probing the MSSM Higgs sector via weak boson fusion at the LHC," *Phys. Lett.* **B454** (1999) 297–303, arXiv:hep-ph/9902434.
- [481] C. Foudas, A. Nikitenko, and M. Takahashi, "Observation of the Standard Model Higgs boson via $H \rightarrow \tau\tau \rightarrow \text{lepton+jet}$ Channel," *CMS Note* **2006/088** (2006).
- [482] H. Pi, P. Avery, J. Rohlf, C. Tully, and S. Kunori, "Search for Standard Model Higgs Boson via Vector Boson Fusion in the $H \rightarrow W^+W^- \rightarrow \ell^\pm \nu jj$ with $120 < m_H < 250$ GeV/ c^2 ," *CMS Note* **2006/092** (2006).
- [483] M. Dubinin, V. Litvin, Y. Ma, H. Newman, and M. Pieri, "Vector Boson Fusion Production with $H \rightarrow \gamma\gamma$," *CMS Note* **2006/097** (2006).
- [484] C. Delaere, "Study of associated WH production with $H \rightarrow WW^*$ in the 3 leptons final state.," *CMS Note* **2006/053** (2006).
- [485] CDF Collaboration, D. Acosta et al., "Measurement of the $t\bar{t}$ Production Cross Section in $pp\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV using Lepton + Jets Events with Secondary Vertex b-tagging," *Phys. Rev.* **D71** (2005) 052003.
- [486] A. Djouadi and S. Ferrag, "PDF Uncertainties In Higgs Production At Hadron Colliders," arXiv:hep-ph/0310209.
- [487] Higgs Working Group Collaboration, K. Assamagan et al., "The Higgs working group: Summary report 2003," arXiv:hep-ph/0406152.
- [488] G. Belanger, F. Boudjema, and K. Sridhar, "SUSY Higgs at the LHC: Large stop mixing effects and associated production," *Nucl. Phys.* **B568** (2000) 3–39, arXiv:hep-ph/9904348.
- [489] M. Dubinin, V. Ilyin, and V. Savrin, "Light Higgs Boson Signal at LHC in the reaction $pp \rightarrow \gamma\gamma + \text{jet}$ and $pp \rightarrow \gamma\gamma + \text{lepton}$," *CMS Note* **1997/101** (1997).
- [490] R. Kinnunen and D. Denegri, "Expected SM/SUSY Higgs Observability in CMS," *CMS Note* **1997/057** (1997).
- [491] ATLAS Collaboration, "ATLAS Detector and Physics Performance. Technical Design Report. Vol. 2," CERN/LHCC CERN-LHCC-99-15 (1999).
- [492] P. Beauchemin, G. Azuelos, and C. Burgess, "Dimensionless coupling of bulk scalars at the LHC," *J. Phys.* **G30** (2004) N17. doi:10.1088/0954-3899/30/10/N01.
- [493] T. Stelzer and W. Long, "Automatic generation of tree level helicity amplitudes," *Comput. Phys. Commun.* **81** (1994) 357–371. doi:10.1016/0010-4655(94)90084-1.
- [494] H. Murayama, I. Watanabe, and K. Hagiwara, "HELAS: HELicity amplitude subroutines for Feynman diagram evaluations." KEK-91-11.
- [495] M. Mangano, M. Moretti, and R. Pittau, "Multijet matrix elements and shower evolution in hadronic collisions: $Wb\bar{b} + n\text{-jets}$ as a case study," *Nucl. Phys.* **B632** (2002) 343–362, arXiv:hep-ph/0108069. doi:10.1016/S0550-3213(02)00249-3.

- [496] F. Caravaglios, M. Mangano, M. Moretti, and R. Pittau, "A new approach to multi-jet calculations in hadron collisions," *Nucl. Phys.* **B539** (1999) 215–232, arXiv:hep-ph/9807570.
- [497] C. Buttar et al., "Les Houches physics at TeV colliders 2005, standard model, QCD, EW, and Higgs working group: Summary report," 2006.
- [498] S. Bityukov, S. Erofeeva, N. Krasnikov, and A. Nikitenko, "Program for evaluation of significance, confidence intervals and limits by direct calculation of probabilities," *Proceedings of PhyStat 2005* (2005).
- [499] M. Lethuillier et al., "Search for a neutral Higgs boson with WH / ZH, $H \rightarrow \gamma\gamma$ channel," *CMS Note* **2006/110** (2006).
- [500] O. Ravat, "Etude du Calorimètre électromagnétique de l'expérience CMS et recherche de bosons de Higgs neutres dans le canal de production associée". PhD thesis, IPN, Lyon, 2004. Thesis LYCEN-T2004-29.
- [501] D. Graham, "An algorithm using tracks to locate the two photon vertex at high luminosity," *CMS TN* **1995/115** (1995).
- [502] A. Djouadi, "Squark effects on Higgs boson production and decay at the LHC," *Phys. Lett.* **B435** (1998) 101–108, arXiv:hep-ph/9806315.
- [503] G. Eynard, "Study of associated Higgs boson production $HW, H\bar{t}\bar{t}, HZ \rightarrow \gamma\gamma + e^\pm/\mu^\pm + X$ with the ATLAS detector at LHC. (In French)". PhD thesis, Annecy, 1998. CERN-THESIS-2000-036.
- [504] P.-H. Beauchemin and G. Azuelos, "Search for the Standard Model Higgs Boson in the $\gamma\gamma + E_T^{miss}$ channel," *ATL-PHYS-2004-028* (2004).
- [505] M. Lethuillier, J.-L. Agram, C. Baty, S. Gascon-Shotkin, S. Perries, and O. Ravat, "Search for a Neutral Higgs Boson with WH/ZH, $H \rightarrow \gamma\gamma$ Channel," *CMS Note* **2006/110** (2006).
- [506] CMS Collaboration, "The TriDAS Project Technical Design Report, Volume 1: The Trigger Systems," *CERN/LHCC* **2000-38** (2000). CMS TDR 6.1.
- [507] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume 1, Section 10.3.2: Photon isolation," *CERN/LHCC* **2006-001** (2006) 376.
- [508] A. L. Read, "Presentation of search results: The CL(s) technique," *J. Phys.* **G28** (2002) 2693–2704.
- [509] T. Junk, "Confidence level computation for combining searches with small statistics," *Nucl. Instrum. Meth.* **A434** (1999) 435–443, arXiv:hep-ex/9902006.
- [510] A. Djouadi and S. Ferrag, "PDF uncertainties in Higgs production at hadron colliders," *Phys. Lett.* **B586** (2004) 345–352, arXiv:hep-ph/0310209.
- [511] J. R. Dell'Aquila and C. A. Nelson, "Simple tests for CP or P violation by sequential decays: $V_1 V_2$ modes with decays into $\bar{\ell}_A \ell_B$ and/or $\bar{q}_A q_B$," *Phys. Rev.* **D33** (1986) 101. doi:10.1103/PhysRevD.33.101.

- [512] A. Skjold and P. Osland, "Signals of CP violation in Higgs decay," *Phys. Lett.* **B329** (1994) 305–311, arXiv:hep-ph/9402358.
- [513] S. Y. Choi, D. J. Miller, M. M. Muhlleitner, and P. M. Zerwas, "Identifying the Higgs spin and parity in decays to Z pairs," *Phys. Lett.* **B553** (2003) 61–71, arXiv:hep-ph/0210077.
- [514] C. P. Buszello, I. Fleck, P. Marquard, and J. J. van der Bij, "Prospective analysis of spin- and CP-sensitive variables in $H \rightarrow ZZ \rightarrow l(1)+l(1)-l(2)+l(2)-$ at the LHC," *Eur. Phys. J.* **C32** (2004) 209–219, arXiv:hep-ph/0212396.
- [515] M. Bluj, "A Study of Angular Correlations in $H \rightarrow ZZ \rightarrow 2e2\mu$," *CMS Note* **2006/094** (2006).
- [516] J. Wess and B. Zumino, "Supergauge Transformations in Four-Dimensions," *Nucl. Phys.* **B70** (1974) 39–50.
- [517] P. Fayet and S. Ferrara, "Supersymmetry," *Phys. Rept.* **32** (1977) 249–334.
- [518] H. P. Nilles, "Supersymmetry, Supergravity and Particle Physics," *Phys. Rept.* **110** (1984) 1.
- [519] R. Barbieri, "Looking Beyond the Standard Model: the Supersymmetric Option," *Riv. Nuovo Cim.* **11N4** (1988) 1–45.
- [520] H. E. Haber and G. L. Kane, "The Search for Supersymmetry: Probing Physics Beyond the Standard Model," *Phys. Rept.* **117** (1985) 75.
- [521] E. Witten, "Mass Hierarchies in Supersymmetric Theories," *Phys. Lett.* **B105** (1981) 267.
- [522] S. Dimopoulos, S. Raby, and F. Wilczek, "Supersymmetry and the Scale of Unification," *Phys. Rev.* **D24** (1981) 1681–1683.
- [523] L. E. Ibanez and G. G. Ross, "Low-Energy Predictions in Supersymmetric Grand Unified Theories," *Phys. Lett.* **B105** (1981) 439.
- [524] L. E. Ibanez and G. G. Ross, " $SU(2)_L \times U(1)$ Symmetry Breaking as a Radiative Effect of Supersymmetry Breaking in GUTs," *Phys. Lett.* **B110** (1982) 215–220. doi:10.1016/0370-2693(82)91239-4.
- [525] K. Inoue, A. Kakuto, H. Komatsu, and S. Takeshita, "Aspects of Grand Unified Models with Softly Broken Supersymmetry," *Prog. Theor. Phys.* **68** (1982) 927.
- [526] L. Alvarez-Gaume, M. Claudson, and M. B. Wise, "Low-Energy Supersymmetry," *Nucl. Phys.* **B207** (1982) 96.
- [527] J. R. Ellis, D. V. Nanopoulos, and K. Tamvakis, "Grand Unification in Simple Supergravity," *Phys. Lett.* **B121** (1983) 123.
- [528] S. Dimopoulos and H. Georgi, "Softly Broken Supersymmetry and $SU(5)$," *Nucl. Phys.* **B193** (1981) 150.
- [529] N. Sakai, "Naturalness in Supersymmetric 'GUTS'," *Zeit. Phys.* **C11** (1981) 153.

- [530] Y. Okada, M. Yamaguchi, and T. Yanagida, "Upper bound of the lightest Higgs boson mass in the minimal supersymmetric standard model," *Prog. Theor. Phys.* **85** (1991) 1–6.
- [531] H. E. Haber and R. Hempfling, "Can the mass of the lightest Higgs boson of the minimal supersymmetric model be larger than $m(Z)$?", *Phys. Rev. Lett.* **66** (1991) 1815–1818.
- [532] J. R. Ellis, G. Ridolfi, and F. Zwirner, "Radiative corrections to the masses of supersymmetric Higgs bosons," *Phys. Lett.* **B257** (1991) 83–91.
- [533] R. Barbieri, M. Frigeni, and F. Caravaglios, "The Supersymmetric Higgs for heavy superpartners," *Phys. Lett.* **B258** (1991) 167–170.
- [534] A. Yamada, "Radiative corrections to the Higgs masses in the minimal supersymmetric standard model," *Phys. Lett.* **B263** (1991) 233–238.
- [535] A. Brignole, J. R. Ellis, G. Ridolfi, and F. Zwirner, "The Supersymmetric charged Higgs boson mass and LEP phenomenology," *Phys. Lett.* **B271** (1991) 123–132.
- [536] P. H. Chankowski, S. Pokorski, and J. Rosiek, "Charged and neutral supersymmetric Higgs boson masses: Complete one loop analysis," *Phys. Lett.* **B274** (1992) 191–198.
- [537] J. Espinosa and M. Quiros, "Two loop radiative corrections to the mass of the lightest Higgs boson in supersymmetric standard models," *Phys. Lett.* **B266** (1991) 389–396.
- [538] R. Hempfling and A. H. Hoang, "Two loop radiative corrections to the upper limit of the lightest Higgs boson mass in the minimal supersymmetric model," *Phys. Lett.* **B331** (1994) 99–106, [arXiv:hep-ph/9401219](https://arxiv.org/abs/hep-ph/9401219).
- [539] J. Casas, J. Espinosa, M. Quiros, and A. Riotto, "The Lightest Higgs boson mass in the minimal supersymmetric standard model," *Nucl. Phys.* **B436** (1995) 3–29, [arXiv:hep-ph/9407389](https://arxiv.org/abs/hep-ph/9407389).
- [540] M. Carena, J. Espinosa, M. Quiros, and C. Wagner, "Analytical expressions for radiatively corrected Higgs masses and couplings in the MSSM," *Phys. Lett.* **B355** (1995) 209–221, [arXiv:hep-ph/9504316](https://arxiv.org/abs/hep-ph/9504316).
- [541] M. Carena, M. Quiros, and C. Wagner, "Effective potential methods and the Higgs mass spectrum in the MSSM," *Nucl. Phys.* **B461** (1996) 407–436, [arXiv:hep-ph/9508343](https://arxiv.org/abs/hep-ph/9508343).
- [542] S. Heinemeyer, W. Hollik, and G. Weiglein, "QCD corrections to the masses of the neutral CP-even Higgs bosons in the MSSM," *Phys. Rev.* **D58** (1998) 091701, [arXiv:hep-ph/9803277](https://arxiv.org/abs/hep-ph/9803277).
- [543] S. Heinemeyer, W. Hollik, and G. Weiglein, "Precise prediction for the mass of the lightest Higgs boson in the MSSM," *Phys. Lett.* **B440** (1998) 296–304, [arXiv:hep-ph/9807423](https://arxiv.org/abs/hep-ph/9807423).
- [544] S. Heinemeyer, W. Hollik, and G. Weiglein, "Constraints on $\tan(\beta)$ in the MSSM from the upper bound on the mass of the lightest Higgs boson," *JHEP* **06** (2000) 009, [arXiv:hep-ph/9909540](https://arxiv.org/abs/hep-ph/9909540).

- [545] J. R. Espinosa and R.-J. Zhang, "Complete two-loop dominant corrections to the mass of the lightest CP-even Higgs boson in the minimal supersymmetric standard model," *Nucl. Phys.* **B586** (2000) 3–38, arXiv:hep-ph/0003246.
- [546] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, "On the $\mathcal{O}(\alpha_t^2)$ two-loop corrections to the neutral Higgs boson masses in the MSSM," *Nucl. Phys.* **B631** (2002) 195–218. doi:10.1016/S0550-3213(02)00184-0.
- [547] A. Brignole, G. Degrassi, P. Slavich, and F. Zwirner, "On the two-loop sbottom corrections to the neutral Higgs boson masses in the MSSM," *Nucl. Phys.* **B643** (2002) 79–92, arXiv:hep-ph/0206101.
- [548] S. Heinemeyer, W. Hollik, H. Rzehak, and G. Weiglein, "High-precision predictions for the MSSM Higgs sector at $\mathcal{O}(\alpha_b\alpha_s)$," *Eur. Phys. J.* **C39** (2005) 465–481. doi:10.1140/epjc/s2005-02112-6.
- [549] J. A. Coarasa, R. A. Jimenez, and J. Sola, "Strong effects on the hadronic widths of the neutral Higgs Bosons in the MSSM," *Phys. Lett.* **B389** (1996) 312–320, arXiv:hep-ph/9511402.
- [550] P. H. Chankowski, S. Pokorski, and J. Rosiek, "Complete on-shell renormalization scheme for the minimal supersymmetric Higgs sector," *Nucl. Phys.* **B423** (1994) 437–496, arXiv:hep-ph/9303309.
- [551] A. Dabelstein, "The One loop renormalization of the MSSM Higgs sector and its application to the neutral scalar Higgs masses," *Z. Phys.* **C67** (1995) 495–512, arXiv:hep-ph/9409375.
- [552] A. Brignole, "Radiative corrections to the supersymmetric neutral Higgs boson masses," *Phys. Lett.* **B281** (1992) 284–294.
- [553] H. E. Haber, R. Hempfling, and A. H. Hoang, "Approximating the radiatively corrected Higgs mass in the minimal supersymmetric model," *Z. Phys.* **C75** (1997) 539–554, arXiv:hep-ph/9609331.
- [554] R.-J. Zhang, "Two-loop effective potential calculation of the lightest CP-even Higgs-boson mass in the MSSM," *Phys. Lett.* **B447** (1999) 89–97, arXiv:hep-ph/9808299.
- [555] J. R. Espinosa and R.-J. Zhang, "MSSM lightest CP-even Higgs boson mass to $\mathcal{O}(\alpha(s)\alpha(t))$: The effective potential approach," *JHEP* **03** (2000) 026, arXiv:hep-ph/9912236.
- [556] G. Degrassi, P. Slavich, and F. Zwirner, "On the neutral Higgs boson masses in the MSSM for arbitrary stop mixing," *Nucl. Phys.* **B611** (2001) 403–422, arXiv:hep-ph/0105096.
- [557] R. Hempfling, "Yukawa coupling unification with supersymmetric threshold corrections," *Phys. Rev.* **D49** (1994) 6168–6172.
- [558] L. J. Hall, R. Rattazzi, and U. Sarid, "The Top quark mass in supersymmetric SO(10) unification," *Phys. Rev.* **D50** (1994) 7048–7065, arXiv:hep-ph/9306309.

- [559] M. Carena, M. Olechowski, S. Pokorski, and C. Wagner, "Electroweak symmetry breaking and bottom-top Yukawa unification," *Nucl. Phys.* **B426** (1994) 269–300, arXiv:hep-ph/9402253.
- [560] J. Guasch, P. Hafliger, and M. Spira, "MSSM Higgs decays to bottom quark pairs revisited," *Phys. Rev.* **D68** (2003) 115001, arXiv:hep-ph/0305101.
- [561] A. Dedes, G. Degrassi, and P. Slavich, "On the two-loop Yukawa corrections to the MSSM Higgs boson masses at large $\tan(\beta)$," *Nucl. Phys.* **B672** (2003) 144–162, arXiv:hep-ph/0305127.
- [562] H. Eberl, K. Hidaka, S. Kraml, W. Majerotto, and Y. Yamada, "Improved SUSY QCD corrections to Higgs boson decays into quarks and squarks," *Phys. Rev.* **D62** (2000) 055006, arXiv:hep-ph/9912463.
- [563] M. Carena, D. Garcia, U. Nierste, and C. E. Wagner, "Effective Lagrangian for the $\bar{t}bH^+$ interaction in the MSSM and charged Higgs phenomenology," *Nucl. Phys.* **B577** (2000) 88–120. doi:0.1016/S0550-3213(00)00146-2.
- [564] S. Heinemeyer, W. Hollik, and G. Weiglein, "Electroweak precision observables in the minimal supersymmetric standard model," *Phys. Rept.* **425** (2006) 265–368, arXiv:hep-ph/0412214.
- [565] M. Carena and H. E. Haber, "Higgs boson theory and phenomenology. ((V))," *Prog. Part. Nucl. Phys.* **50** (2003) 63–152, arXiv:hep-ph/0208209.
- [566] **ALEPH, DELPHI, L3, OPAL** Collaboration, LEP Working Group for Higgs Boson Searches, "Search for neutral MSSM Higgs bosons at LEP," arXiv:hep-ex/0602042.
- [567] **LEP Higgs Working Group for Higgs boson searches** Collaboration, "Search for charged Higgs bosons: Preliminary combined results using LEP data collected at energies up to 209- GeV," arXiv:hep-ex/0107031.
- [568] A. Dabelstein, "Fermionic decays of neutral MSSM Higgs bosons at the one loop level," *Nucl. Phys.* **B456** (1995) 25–56, arXiv:hep-ph/9503443.
- [569] S. Heinemeyer, W. Hollik, and G. Weiglein, "Decay widths of the neutral CP-even MSSM Higgs bosons in the Feynman-diagrammatic approach," *Eur. Phys. J.* **C16** (2000) 139–153, arXiv:hep-ph/0003022.
- [570] A. Djouadi, P. Janot, J. Kalinowski, and P. M. Zerwas, "SUSY Decays of Higgs Particles," *Phys. Lett.* **B376** (1996) 220–226, arXiv:hep-ph/9603368.
- [571] A. Djouadi, J. Kalinowski, P. Ohmann, and P. M. Zerwas, "Heavy SUSY Higgs bosons at e^+e^- linear colliders," *Z. Phys.* **C74** (1997) 93–111, arXiv:hep-ph/9605339.
- [572] S. Dawson, A. Djouadi, and M. Spira, "QCD Corrections to SUSY Higgs Production: The Role of Squark Loops," *Phys. Rev. Lett.* **77** (1996) 16–19, arXiv:hep-ph/9603423.
- [573] R. V. Harlander and M. Steinhauser, "Hadronic Higgs production and decay in supersymmetry at next-to-leading order," *Phys. Lett.* **B574** (2003) 258–268, arXiv:hep-ph/0307346.

- [574] R. Harlander and M. Steinhauser, "Effects of SUSY-QCD in hadronic Higgs production at next- to-next-to-leading order," *Phys. Rev.* **D68** (2003) 111701, arXiv:hep-ph/0308210.
- [575] R. V. Harlander and M. Steinhauser, "Supersymmetric Higgs production in gluon fusion at next-to- leading order," *JHEP* **09** (2004) 066, arXiv:hep-ph/0409010.
- [576] R. V. Harlander and F. Hofmann, "Pseudo-scalar Higgs production at next-to-leading order SUSY-QCD," arXiv:hep-ph/0507041.
- [577] U. Langenegger, M. Spira, A. Starodumov, and P. Trueb, "SM and MSSM Higgs boson production: Spectra at large transverse momentum," arXiv:hep-ph/0604156.
- [578] A. Djouadi and M. Spira, "SUSY-QCD corrections to Higgs boson production at hadron colliders," *Phys. Rev.* **D62** (2000) 014004, arXiv:hep-ph/9912476.
- [579] P. Wu et al., "NLO supersymmetric QCD corrections to t anti-t h0 associated production at hadron colliders," arXiv:hep-ph/0505086.
- [580] S. Dittmaier, M. Kramer, and M. Spira, "Higgs radiation off bottom quarks at the Tevatron and the LHC," *Phys. Rev.* **D70** (2004) 074010, arXiv:hep-ph/0309204.
- [581] S. Dawson, C. B. Jackson, L. Reina, and D. Wackerroth, "Exclusive Higgs boson production with bottom quarks at hadron colliders," *Phys. Rev.* **D69** (2004) 074027, arXiv:hep-ph/0311067.
- [582] D. A. Dicus and S. Willenbrock, "Higgs Boson Production from Heavy Quark Fusion," *Phys. Rev.* **D39** (1989) 751.
- [583] D. Dicus, T. Stelzer, Z. Sullivan, and S. Willenbrock, "Higgs boson production in association with bottom quarks at next-to-leading order," *Phys. Rev.* **D59** (1999) 094016, arXiv:hep-ph/9811492.
- [584] C. Balazs, H.-J. He, and C. P. Yuan, "QCD corrections to scalar production via heavy quark fusion at hadron colliders," *Phys. Rev.* **D60** (1999) 114001, arXiv:hep-ph/9812263.
- [585] R. V. Harlander and W. B. Kilgore, "Higgs boson production in bottom quark fusion at next-to- next-to-leading order," *Phys. Rev.* **D68** (2003) 013001, arXiv:hep-ph/0304035.
- [586] M. A. G. Aivazis, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 1. General formalism and kinematics of charged current and neutral current production processes," *Phys. Rev.* **D50** (1994) 3085–3101, arXiv:hep-ph/9312318.
- [587] M. A. G. Aivazis, J. C. Collins, F. I. Olness, and W.-K. Tung, "Leptoproduction of heavy quarks. 2. A Unified QCD formulation of charged and neutral current processes from fixed target to collider energies," *Phys. Rev.* **D50** (1994) 3102–3118, arXiv:hep-ph/9312319.
- [588] J. Campbell et al., "Higgs boson production in association with bottom quarks," arXiv:hep-ph/0405302. Contributed to 3rd Les Houches Workshop: Physics at TeV Colliders, Les Houches, France, 26 May - 6 Jun 2003.

- [589] J. Campbell, R. K. Ellis, F. Maltoni, and S. Willenbrock, "Higgs boson production in association with a single bottom quark," *Phys. Rev.* **D67** (2003) 095002, arXiv:hep-ph/0204093.
- [590] S. Dawson, C. B. Jackson, L. Reina, and D. Wackerath, "Higgs boson production with one bottom quark jet at hadron colliders," *Phys. Rev. Lett.* **94** (2005) 031802. doi:10.1103/PhysRevLett.94.031802.
- [591] A. C. Bawa, C. S. Kim, and A. D. Martin, "Charged Higgs Production at Hadron Colliders," *Z. Phys.* **C47** (1990) 75–82.
- [592] F. Borzumati, J.-L. Kneur, and N. Polonsky, "Higgs-strahlung and R-parity violating slepton-strahlung at hadron colliders," *Phys. Rev.* **D60** (1999) 115011, arXiv:hep-ph/9905443.
- [593] A. Belyaev, D. Garcia, J. Guasch, and J. Sola, "Prospects for heavy supersymmetric charged Higgs boson searches at hadron colliders," *JHEP* **06** (2002) 059, arXiv:hep-ph/0203031.
- [594] P. Wu et al., "NLO supersymmetric QCD corrections to the t anti-b H- associated production at hadron colliders," *Phys. Rev.* **D73** (2006) 015012, arXiv:hep-ph/0601069. doi:10.1103/PhysRevD.73.015012.
- [595] S.-H. Zhu, "Complete next-to-leading order QCD corrections to charged Higgs boson associated production with top quark at the CERN Large Hadron Collider," *Phys. Rev.* **D67** (2003) 075006. doi:10.1103/PhysRevD.67.075006.
- [596] T. Plehn, "Charged Higgs boson production in bottom-gluon fusion." Prepared for 10th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY02), Hamburg, Germany, 17-23 Jun 2002.
- [597] E. L. Berger, T. Han, J. Jiang, and T. Plehn, "Associated production of a top quark and a charged Higgs boson," *Phys. Rev.* **D71** (2005) 115012, arXiv:hep-ph/0312286.
- [598] G. Gao, G. Lu, Z. Xiong, and J. M. Yang, "Loop effects and non-decoupling property of SUSY QCD in $gb \rightarrow tH^-$," *Phys. Rev.* **D66** (2002) 015007. doi:10.1103/PhysRevD.66.015007.
- [599] S. S. D. Willenbrock, "Pair Production of Supersymmetric Charged Higgs Bosons," *Phys. Rev.* **D35** (1987) 173.
- [600] A. Krause, T. Plehn, M. Spira, and P. M. Zerwas, "Production of charged Higgs boson pairs in gluon gluon collisions," *Nucl. Phys.* **B519** (1998) 85–100, arXiv:hep-ph/9707430.
- [601] Y. Jiang, L. Han, W.-G. Ma, Z.-H. Yu, and M. Han, "Pair production of charged Higgs bosons in gluon gluon collisions," *J. Phys.* **G23** (1997) 385–400, arXiv:hep-ph/9703275.
- [602] O. Brein and W. Hollik, "Pair production of charged MSSM Higgs bosons by gluon fusion," *Eur. Phys. J.* **C13** (2000) 175–184, arXiv:hep-ph/9908529.

- [603] A. A. Barrientos Bendezu and B. A. Kniehl, “ H^+H^- pair production at the Large Hadron Collider,” *Nucl. Phys.* **B568** (2000) 305–318.
doi:10.1016/S0550-3213(99)00732-4.
- [604] H.-S. Hou et al., “Pair production of charged Higgs bosons from bottom-quark fusion,” *Phys. Rev.* **D71** (2005) 075014. doi:10.1103/PhysRevD.71.075014.
- [605] A. A. Barrientos Bendezu and B. A. Kniehl, “ $W^\pm H^\mp$ associated production at the Large Hadron Collider,” *Phys. Rev.* **D59** (1999) 015009, arXiv:hep-ph/9807480.
doi:10.1103/PhysRevD.59.015009.
- [606] O. Brein, W. Hollik, and S. Kanemura, “The MSSM prediction for $W^\pm H^\mp$ production by gluon fusion,” *Phys. Rev.* **D63** (2001) 095001, arXiv:hep-ph/0008308.
- [607] A. A. Barrientos Bendezu and B. A. Kniehl, “Squark loop correction to $W^{+-}H^{+-}$ associated hadroproduction,” *Phys. Rev.* **D63** (2001) 015009,
arXiv:hep-ph/0007336.
- [608] D. Dicus, J. Hewett, C. Kao, and T. Rizzo, “ $W^\pm H^\mp$ production at hadron colliders,” *Phys. Rev.* **D40** (1989) 787.
- [609] W. Hollik and S.-h. Zhu, “ $\mathcal{O}(\alpha_s)$ corrections to $b\bar{b} \rightarrow W^\pm H^\mp$ at the CERN Large Hadron Collider,” *Phys. Rev.* **D65** (2002) 075015, arXiv:hep-ph/0109103.
doi:10.1103/PhysRevD.65.075015.
- [610] J. Zhao, C. S. Li, and Q. Li, “SUSY-QCD corrections to $W^\pm H^\mp$ associated production at the CERN Large Hadron Collider,” *Phys. Rev.* **D72** (2005) 114008,
arXiv:hep-ph/0509369.
- [611] S. Lehti, “Study of $H/A \rightarrow \tau\tau \rightarrow e\mu + X$ in CMS,” *CMS Note* **2006/101** (2006).
- [612] S. Lehti, “Study of $gg \rightarrow b\bar{b}H_{SUSY}, H_{SUSY} \rightarrow \tau\tau \rightarrow \ell\ell + X$,” *CMS Note* **2002/035** (2002).
- [613] D. Bloch. Private communication.
- [614] E. Boos, A. Djouadi, M. Muhlleitner, and A. Vologdin, “The MSSM Higgs bosons in the intense-coupling regime,” *Phys. Rev.* **D66** (2002) 055004,
arXiv:hep-ph/0205160.
- [615] E. Boos, A. Djouadi, and A. Nikitenko, “Detection of the neutral MSSM Higgs bosons in the intense-coupling regime at the LHC,” *Phys. Lett.* **B578** (2004) 384–393,
arXiv:hep-ph/0307079.
- [616] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 12.2.3: Combined secondary vertex tag,” *CERN/LHCC* **2006-001** (2006) 466.
- [617] J. Fernandez, “Search for MSSM heavy neutral Higgs bosons in the four-b final state,” *CMS Note* **2006/080** (2006).
- [618] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 12.2.6: HLT b tag,” *CERN/LHCC* **2006-001** (2006) 474.
- [619] CMS Collaboration, D. Acosta et al., “CMS Physics TDR Volume 1, Section 11.6.4: Parton-level corrections,” *CERN/LHCC* **2006-001** (2006) 428.

- [620] DØ Collaboration, V. M. Abazov et al., “Search for neutral supersymmetric Higgs bosons in multijet events at $\sqrt{s} = 1.96$ TeV,” *Phys. Rev. Lett.* **95** (2005) 151801. doi:10.1103/PhysRevLett.95.151801.
- [621] M. Baarmand, M. Hashemi, and A. Nikitenko, “Light Charged Higgs Discovery Potential of CMS in the $H^+ \rightarrow \tau\nu$ with Single Lepton Trigger,” *CMS Note* **2006/056** (2006).
- [622] D. Roy, “The hadronic tau decay signature of a heavy charged Higgs boson at LHC,” *Phys. Lett.* **B459** (1999) 607–614, arXiv:hep-ph/9905542.
- [623] R. Kinnunen, “Study of $A/H \rightarrow \tau\tau$ and $H^\pm \rightarrow \tau\nu$ in CMS,” *Czech. J. Phys.* **54** (2004) A93–A101.
- [624] K. A. Assamagan and Y. Coadou, “The hadronic tau decay of a heavy H^\pm in ATLAS,” *Acta Phys. Polon.* **B33** (2002) 707–720.
- [625] M. Baarmand, M. Hashemi, and A. Nikitenko, “Search for the Heavy Charged MSSM Higgs bosons with the $H^+ \rightarrow \tau\nu$ Decay Mode in Fully Hadronic Final State,” *CMS Note* **2006/100** (2006).
- [626] D. Roy, “Looking for the charged Higgs boson,” *Mod. Phys. Lett.* **A19** (2004) 1813–1828, arXiv:hep-ph/0406102.
- [627] J. Alwall and J. Rathsmann, “Improved description of charged Higgs boson production at hadron colliders,” *JHEP* **12** (2004) 050, arXiv:hep-ph/0409094.
- [628] T. Plehn, “Charged Higgs boson production in bottom gluon fusion,” *Phys. Rev.* **D67** (2003) 014018, arXiv:hep-ph/0206121.
- [629] S. Lowette, J. D’Hondt, and P. Vanlaer, “Charged MSSM Higgs Boson Observability in the $H^\pm \rightarrow tb$ Decay,” *CMS Note* **2006/109** (2006).
- [630] CMS Collaboration, G. Anagnostou and G. Daskalakis, “Search for the MSSM $A \rightarrow Zh$ decay with $Z \rightarrow \ell^+\ell^-$, $h \rightarrow b\bar{b}$,” *CMS Note* **2006/063** (2006).
- [631] CMS Collaboration, C. Charlot, R. Salerno, and Y. Sirois, “Observability of the heavy neutral SUSY Higgs Bosons decaying into neutralinos,” *CMS Note* **2006/125** (2006).
- [632] B. C. Allanach et al., “The Snowmass points and slopes: Benchmarks for SUSY searches,” *Eur. Phys. J.* **C25** (2002) 113–123, arXiv:hep-ph/0202233.
- [633] M. Battaglia et al., “Updated post-WMAP benchmarks for supersymmetry,” *Eur. Phys. J.* **C33** (2004) 273–296, arXiv:hep-ph/0306219.
- [634] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, “Suggestions for improved benchmark scenarios for Higgs-boson searches at LEP2,” arXiv:hep-ph/9912223.
- [635] M. Carena, S. Heinemeyer, C. Wagner, and G. Weiglein, “Suggestions for benchmark scenarios for MSSM Higgs boson searches at hadron colliders,” *Eur. Phys. J.* **C26** (2003) 601–607, arXiv:hep-ph/0202167.
- [636] ECFA/DESY LC Physics Working Group Collaboration, J. A. Aguilar-Saavedra et al., “TESLA Technical Design Report Part III: Physics at an e+e- Linear Collider,” arXiv:hep-ph/0106315.

- [637] C. L. Bennett et al., "First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results," *Astrophys. J. Suppl.* **148** (2003) 1, arXiv:astro-ph/0302207.
- [638] WMAP Collaboration, D. N. Spergel et al., "First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters," *Astrophys. J. Suppl.* **148** (2003) 175, arXiv:astro-ph/0302209.
- [639] H. Goldberg, "Constraint on the photino mass from cosmology," *Phys. Rev. Lett.* **50** (1983) 1419.
- [640] J. R. Ellis, J. Hagelin, D. Nanopoulos, K. A. Olive, and M. Srednicki, "Supersymmetric relics from the big bang," *Nucl. Phys.* **B238** (1984) 453–476.
- [641] Heavy Flavor Averaging Group Collaboration, E. Barberio et al., "Averages of b-hadron properties at the end of 2005," arXiv:hep-ex/0603003.
- [642] H. M. Asatrian, A. Hovhannisyan, V. Poghosyan, C. Greub, and T. Hurth, "Towards the NNLL precision in anti-B \rightarrow X/s gamma," arXiv:hep-ph/0512097.
- [643] Muon Collaboration, G. W. Bennett, "Final report of the muon E821 anomalous magnetic moment measurement at BNL," arXiv:hep-ex/0602035.
- [644] A. Czarnecki and W. J. Marciano, "The muon anomalous magnetic moment: A harbinger for 'new physics'," *Phys. Rev.* **D64** (2001) 013014, arXiv:hep-ph/0102122.
- [645] M. Carena et al., "Reconciling the two-loop diagrammatic and effective field theory computations of the mass of the lightest CP-even Higgs boson in the MSSM," *Nucl. Phys.* **B580** (2000) 29–57, arXiv:hep-ph/0001002.
- [646] L. Randall and R. Sundrum, "An alternative to compactification," *Phys. Rev. Lett.* **83** (1999) 4690–4693. doi:10.1103/PhysRevLett.83.4690.
- [647] G. F. Giudice, R. Rattazzi, and J. D. Wells, "Graviscalars from higher-dimensional metrics and curvature- Higgs mixing," *Nucl. Phys.* **B595** (2001) 250–276, arXiv:hep-ph/0002178.
- [648] M. Chaichian, A. Datta, K. Huitu, and Z.-h. Yu, "Radion and Higgs mixing at the LHC," *Phys. Lett.* **B524** (2002) 161–169, arXiv:hep-ph/0110035.
- [649] J. L. Hewett and T. G. Rizzo, "Shifts in the properties of the Higgs boson from radion mixing," *JHEP* **08** (2003) 028, arXiv:hep-ph/0202155.
- [650] D. Dominici, B. Grzadkowski, J. F. Gunion, and M. Toharia, "The scalar sector of the Randall-Sundrum model," *Nucl. Phys.* **B671** (2003) 243–292, arXiv:hep-ph/0206192.
- [651] M. Battaglia, S. De Curtis, A. De Roeck, D. Dominici, and J. F. Gunion, "On the complementarity of Higgs and radion searches at LHC," *Phys. Lett.* **B568** (2003) 92–102, arXiv:hep-ph/0304245.
- [652] G. Azuelos, D. Cavalli, H. Przysiezniak, and L. Vacavant, "Search for the radion using the ATLAS detector," *Eur. Phys. J. direct* **C4** (2002) 16.

- [653] D. Dominici, G. Dewhurst, A. Nikitenko, S. Gennai, and L. Fanò, "Search for radion decays into Higgs boson pairs in the $\gamma\gamma b\bar{b}$, $\tau\tau b\bar{b}$ and $b\bar{b}b\bar{b}$ final states," *CMS Note* **2005/007** (2005).
- [654] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, "(De)constructing dimensions," *Phys. Rev. Lett.* **86** (2001) 4757–4761, arXiv:hep-th/0104005.
- [655] H.-C. Cheng, C. T. Hill, S. Pokorski, and J. Wang, "The standard model in the latticized bulk," *Phys. Rev.* **D64** (2001) 065007, arXiv:hep-th/0104179.
- [656] N. Arkani-Hamed, A. G. Cohen, and H. Georgi, "Electroweak symmetry breaking from dimensional deconstruction," *Phys. Lett.* **B513** (2001) 232–240, arXiv:hep-ph/0105239.
- [657] N. Arkani-Hamed, A. G. Cohen, E. Katz, and A. E. Nelson, "The littlest Higgs," *JHEP* **07** (2002) 034, arXiv:hep-ph/0206021.
- [658] J. Schechter and J. Valle, "Neutrino masses in $SU(2) \times U(1)$ Theories," *Phys. Rev.* **D22** (1980) 2227. doi:10.1103/PhysRevD.22.2227.
- [659] E. Ma and U. Sarkar, "Neutrino masses and leptogenesis with heavy Higgs triplets," *Phys. Rev. Lett.* **80** (1998) 5716–5719, arXiv:hep-ph/9802445.
- [660] E. Ma, M. Raidal, and U. Sarkar, "Verifiable model of neutrino masses from large extra dimensions," *Phys. Rev. Lett.* **85** (2000) 3769–3772, arXiv:hep-ph/0006046.
- [661] E. Ma, M. Raidal, and U. Sarkar, "Phenomenology of the neutrino-mass-giving Higgs triplet and the low-energy seesaw violation of lepton number," *Nucl. Phys.* **B615** (2001) 313–330, arXiv:hep-ph/0012101.
- [662] E. Ma and U. Sarkar, "Connecting dark energy to neutrinos with an observable Higgs triplet," arXiv:hep-ph/0602116.
- [663] G. Marandella, C. Schappacher, and A. Strumia, "Little-Higgs corrections to precision data after LEP2," *Phys. Rev.* **D72** (2005) 035014, arXiv:hep-ph/0502096.
- [664] K. Huitu, J. Maalampi, A. Pietila, and M. Raidal, "Doubly charged Higgs at LHC," *Nucl. Phys.* **B487** (1997) 27–42, arXiv:hep-ph/9606311.
- [665] T. Han, H. E. Logan, and L.-T. Wang, "Smoking-gun signatures of little Higgs models," arXiv:hep-ph/0506313.
- [666] G. Azuelos et al., "Exploring little Higgs models with ATLAS at the LHC," *Eur. Phys. J.* **C39S2** (2005) 13–24, arXiv:hep-ph/0402037.
- [667] J. F. Gunion, J. Grifols, A. Mendez, B. Kayser, and F. I. Olness, "Higgs bosons in left-right symmetric models," *Phys. Rev.* **D40** (1989) 1546. doi:10.1103/PhysRevD.40.1546.
- [668] M. Muhlleitner and M. Spira, "A note on doubly-charged Higgs pair production at hadron colliders," *Phys. Rev.* **D68** (2003) 117701, arXiv:hep-ph/0305288.
- [669] S. Bitjukov, N. Krasnikov, and V. Taperechkina, "Confidence intervals for Poisson distribution parameter," arXiv:hep-ex/0108020.

- [670] S. P. Martin, "A supersymmetry primer," in *Perspectives in Supersymmetry*, ed. G. Kane, World Scientific (1997) arXiv:hep-ph/9709356.
- [671] CMS Collaboration, S. Abdullin et al., "Discovery potential for supersymmetry in CMS," *J. Phys.* **G28** (2002) 469. doi:10.1088/0954-3899/28/3/401.
- [672] F. E. Paige, S. D. Protopopescu, H. Baer, and X. Tata, "ISAJET 7.69: A Monte Carlo Event Generator for pp , $\bar{p}p$, and e^+e^- Reactions," arXiv:hep-ph/0312045.
- [673] A. De Roeck et al., "Supersymmetric benchmarks with non-universal scalar masses or gravitino dark matter," arXiv:hep-ph/0508198.
- [674] W. de Boer et al., "Excess of EGRET galactic gamma ray data interpreted as dark matter annihilation," arXiv:astro-ph/0408272.
- [675] D. Acosta et al., "Potential to Discover SUSY in Events with Muons Jets and Large Missing Transverse Energy in pp Collisions at $\sqrt{s} = 14$ TeV," *CMS Note* **2006/134** (2006).
- [676] D. Acosta et al., "CMS Discovery Potential for mSUGRA in Same Sign Di-muon Events with Jets and Large Missing Transverse Energy in pp collisions at $\sqrt{s} = 14$ TeV," *CMS Note* (2006).
- [677] M. Chiorboli, M. Galanti, and A. Tricoli, "Leptons + Jets + Missing Energy analysis at LM1," *CMS Note* **2006/133** (2006).
- [678] D. Mangeol and U. Goerlach, "Search for $\tilde{\tau}$ production in di-tau final states and measurements of SUSY masses in mSUGRA cascade decays," *CMS Note* **2006/096** (2006).
- [679] S. Bitjukov, "Uncertainty, Systematics, Limits." Available at <http://cmsdoc.cern.ch/bitjukov>, 2005.
- [680] S. Kyriazopoulou and C. Markou, "Search for SUSY in Final States with Z Bosons," *CMS Note* **2006/116** (2006).
- [681] A. Nikitenko, G. Quast, and V. Ciulli, "Minimal requirements for significance estimates of the physics result." Info located at <http://cmsdoc.cern.ch/~anikiten/cms-higgs/stat.txt>.
- [682] W. Beenakker, R. Hopker, and M. Spira, "PROSPINO: A program for the PROduction of Supersymmetric Particles In Next-to-leading Order QCD," arXiv:hep-ph/9611232.
- [683] B. C. Allanach, C. G. Lester, M. A. Parker, and B. R. Webber, "Measuring sparticle masses in non-universal string inspired models at the LHC," *JHEP* **09** (2000) 004, arXiv:hep-ph/0007009.
- [684] W. Beenakker et al., "The production of charginos/neutralinos and sleptons at hadron colliders," *Phys. Rev. Lett.* **83** (1999) 3780–3783. doi:10.1103/PhysRevLett.83.3780.
- [685] F. del Aguila and L. Ametler, "On the detectability of sleptons at large hadron colliders," *Phys. Lett.* **B261** (1991) 326–333. doi:10.1016/0370-2693(91)90336-0.

- [686] H. Baer, C.-h. Chen, F. Paige, and X. Tata, "Detecting sleptons at hadron colliders and supercolliders," *Phys. Rev.* **D49** (1994) 3283–3290, arXiv:hep-ph/9311248.
- [687] D. Denegri, L. Rurua, and N. Stepanov, "Detection of Sleptons in CMS, Mass Reach," *CMS TN* **96-059** (1996).
- [688] Y. M. Andreev, S. I. Bityukov, and N. V. Krasnikov, "Sleptons at post-WMAP benchmark points at LHC(CMS)," *Phys. Atom. Nucl.* **68** (2005) 340–347, arXiv:hep-ph/0402229.
- [689] S. Bityukov and N. Krasnikov, "The search for sleptons and lepton-flavor-number violation at LHC (CMS)," *Phys. Atom. Nucl.* **62** (1999) 1213–1225, arXiv:hep-ph/9712358.
- [690] N. Krasnikov, "Flavor lepton number violation at LEP-2," *Mod. Phys. Lett.* **A9** (1994) 791–794.
- [691] N. Arkani-Hamed, H.-C. Cheng, J. L. Feng, and L. J. Hall, "Probing Lepton Flavor Violation at Future Colliders," *Phys. Rev. Lett.* **77** (1996) 1937–1940, arXiv:hep-ph/9603431. doi:10.1103/PhysRevLett.77.1937.
- [692] N. Krasnikov, "Search for flavor lepton number violation in slepton decays at LHC," *JETP Lett.* **65** (1997) 148. doi:10.1134/1.567315.
- [693] K. Agashe and M. Graesser, "Signals of supersymmetric lepton flavor violation at the LHC," *Phys. Rev.* **D61** (2000) 075008, arXiv:hep-ph/9904422.
- [694] J. Hisano, R. Kitano, and M. M. Nojiri, "Flavor mixing in slepton production at the large hadron collider," *Phys. Rev.* **D65** (2002) 116002.
- [695] H. Baer, C.-h. Chen, F. Paige, and X. Tata, "Trileptons from chargino - neutralino production at the CERN Large Hadron Collider," *Phys. Rev.* **D50** (1994) 4508–4516, arXiv:hep-ph/9404212. doi:10.1103/PhysRevD.50.4508.
- [696] J. R. Ellis, K. A. Olive, and Y. Santoso, "The MSSM parameter space with non-universal Higgs masses," *Phys. Lett.* **B539** (2002) 107–118, arXiv:hep-ph/0204192.
- [697] G. Altarelli, B. Mele, and M. Ruiz-Altaba, "Searching for New Heavy Vector Bosons in $p\bar{p}$ Colliders," *Z. Phys.* **C45** (1989) 109. doi:10.1007/BF01556677.
- [698] N. Arkani-Hamed, S. Dimopoulos, and G. R. Dvali, "The hierarchy problem and new dimensions at a millimeter," *Phys. Lett.* **B429** (1998) 263–272, arXiv:hep-ph/9803315.
- [699] J. D. Lykken, E. Poppitz, and S. P. Trivedi, "Branes with GUTs and supersymmetry breaking," *Nucl. Phys.* **B543** (1999) 105–121, arXiv:hep-th/9806080.
- [700] I. Antoniadis, "A Possible new dimension at a few TeV," *Phys. Lett.* **B246** (1990) 377–384. doi:10.1016/0370-2693(90)90617-F.
- [701] J. D. Lykken, "Weak Scale Superstrings," *Phys. Rev.* **D54** (1996) 3693–3697, arXiv:hep-th/9603133. doi:10.1103/PhysRevD.54.R3693.

- [702] G. F. Giudice, R. Rattazzi, and J. D. Wells, "Quantum gravity and extra dimensions at high-energy colliders," *Nucl. Phys.* **B544** (1999) 3–38, arXiv:hep-ph/9811291.
- [703] T. Han, J. D. Lykken, and R.-J. Zhang, "On Kaluza-Klein states from large extra dimensions," *Phys. Rev.* **D59** (1999) 105006, arXiv:hep-ph/9811350. doi:10.1103/PhysRevD.59.105006.
- [704] S. Cullen, M. Perelstein, and M. E. Peskin, "TeV strings and collider probes of large extra dimensions," *Phys. Rev.* **D62** (2000) 055012, arXiv:hep-ph/0001166.
- [705] J. D. Lykken and L. Randall, "The shape of gravity," *JHEP* **06** (2000) 014, arXiv:hep-th/9908076.
- [706] L. Vacavant and I. Hinchliffe, "Signals of models with large extra dimensions in ATLAS," *J. Phys.* **G27** (2001) 1839–1850.
- [707] J. Ruppert, C. Rahmede, and M. Bleicher, "Determination of the fundamental scale of gravity and the number of space-time dimensions from high energetic particle interactions," *Phys. Lett.* **B608** (2005) 240–243, arXiv:hep-ph/0501028.
- [708] E. Dudas and J. Mourad, "String theory predictions for future accelerators," *Nucl. Phys.* **B575** (2000) 3–34. doi:10.1016/S0550-3213(00)00082-1.
- [709] D. Chialva, R. Iengo, and J. G. Russo, "Cross sections for production of closed superstrings at high energy colliders in brane world models," *Phys. Rev.* **D71** (2005) 106009, arXiv:hep-ph/0503125.
- [710] M. Bando, T. Kugo, T. Noguchi, and K. Yoshioka, "Brane fluctuation and suppression of Kaluza-Klein mode couplings," *Phys. Rev. Lett.* **83** (1999) 3601–3604. doi:10.1103/PhysRevLett.83.3601.
- [711] J. L. Hewett, "Indirect collider signals for extra dimensions," *Phys. Rev. Lett.* **82** (1999) 4765–4768. doi:10.1103/PhysRevLett.82.4765.
- [712] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, "Phenomenology of the Randall-Sundrum gauge hierarchy model," *Phys. Rev. Lett.* **84** (2000) 2080. doi:10.1103/PhysRevLett.84.2080.
- [713] J. Hewett and M. Spiropulu, "Particle physics probes of extra spacetime dimensions," *Ann. Rev. Nucl. Part. Sci.* **52** (2002) 397–424. doi:10.1146/annurev.nucl.52.050102.090706.
- [714] B. Clerbaux, T. Mahmoud, C. Collard, and P. Miné, "Search with the CMS detector for heavy resonances decaying into an electron pair," *CMS Note* **2006/083** (2006).
- [715] C. Collard and M.-C. Lemaire, "Search with the CMS Detector for Randall-Sundrum Excitations of Gravitons Decaying Into Electron Pairs," *CMS Note* **2004/024** (2004).
- [716] CMS Collaboration, D. Acosta et al., "CMS Physics TDR Volume1, Section 10," *CERN/LHCC* **2006-001** (2006).
- [717] B. Clerbaux, T. Mahmoud, C. Collard, M.-C. Lemaire, and V. Litvin, "TeV electron and photon saturation studies," *CMS Note* **2006/004** (2006).

- [718] CDF Collaboration, T. Affolder et al., "Measurement of ds/dM and Forward-Backward Charge Asymmetry for High-mass Drell-Yan e^+e^- Pairs from collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **87** (2001) 131802.
- [719] CTEQ Collaboration, H. L. Lai et al., "Global QCD analysis of parton structure of the nucleon: CTEQ5 parton distributions," *Eur. Phys. J.* **C12** (2000) 375–392.
doi:10.1007/s100529900196.
- [720] CTEQ Collaboration, D. Stump et al., "Inclusive Jet Production, Parton Distributions, and the Search for New Physics," *JHEP* **0310** (2003) 046, arXiv:hep-ph/0303013.
- [721] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, "Experimental probes of localized gravity: On and off the wall," *Phys. Rev.* **D63** (2001) 075004,
arXiv:hep-ph/0006041.
- [722] K.-m. Cheung and G. Landsberg, "Drell-Yan and diphoton production at hadron colliders and low scale gravity model," *Phys. Rev.* **D62** (2000) 076003,
arXiv:hep-ph/9909218.
- [723] I. Belotelov, I. Golutvin, D. Bourilkov, A. Lanyov, E. Rogalev, M. Savina, and S. Shmatov, "Search for ADD extra dimensional gravity in di-muon channel with the CMS Detector," *CMS Note* **2006/076** (2006).
- [724] R. Mohapatra and J. Pati, "Left-right gauge symmetry and an 'isoconjugate' model of CP violation," *Phys. Rev.* **D11** (1975) 566–571. doi:10.1103/PhysRevD.11.566.
- [725] G. Senjanovic and R. Mohapatra, "Exact Left-Right Symmetry and Spontaneous Violation of Parity," *Phys. Rev.* **D12** (1975) 1502. doi:10.1103/PhysRevD.12.1502.
- [726] R. Mohapatra and G. Senjanovic, "Neutrino mass and spontaneous parity nonconservation," *Phys. Rev. Lett.* **44** (1980) 912.
doi:10.1103/PhysRevLett.44.912.
- [727] J. Pati and A. Salam, "Lepton Number as the Fourth Color," *Phys. Rev.* **D10** (1974) 275–289.
- [728] CDF Collaboration, T. Affolder et al., "Search for quark lepton compositeness and a heavy W' boson using the $e\nu$ channel in p anti- p collisions at $s^{**}(1/2) = 1.8$ -TeV," *Phys. Rev. Lett.* **87** (2001) 231803, arXiv:hep-ex/0107008.
- [729] C. Hof, T. Hebbeker, and K. Hoepfner, "Detection of New Heavy Charged Gauge Bosons in the Muon plus Neutrino Channel," *CMS Note* **2006/117** (2006).
- [730] K. Gumus, N. Akchurin, S. Esen, and R. Harris, "CMS Sensitivity to Dijet Resonances," *CMS Note* **2006/070** (2006).
- [731] U. Baur, I. Hinchliffe, and D. Zeppenfeld, "Excited Quark Production at Hadron Colliders," *Int. J. Mod. Phys.* **A2** (1987) 1285.
- [732] J. Bagger, C. Schmidt, and S. King, "Axigluon Production in Hadronic Collisions," *Phys. Rev.* **D37** (1988) 1188.
- [733] R. S. Chivukula, A. G. Cohen, and E. H. Simmons, "New Strong Interactions at the Tevatron?," *Phys. Lett.* **B380** (1996) 92–98, arXiv:hep-ph/9603311.

- [734] J. L. Hewett and T. G. Rizzo, "Low-Energy Phenomenology of Superstring Inspired E(6) Models," *Phys. Rept.* **183** (1989) 193.
- [735] K. Lane and S. Mrenna, "The collider phenomenology of technihadrons in the technicolor Straw Man Model," *Phys. Rev.* **D67** (2003) 115011.
doi:10.1103/PhysRevD.67.115011.
- [736] E. Eichten, I. Hinchliffe, K. D. Lane, and C. Quigg, "Super Collider Physics," *Rev. Mod. Phys.* **56** (1984) 579–707.
- [737] CMS Collaboration, M.-C. Lemaire, H. Newman, and V. Litvin, "Search for Randall-Sundrum excitations of gravitons decaying into two photons for CMS at LHC," *CMS Note* **2006/051** (2006).
- [738] CMS Collaboration, B. Clerbaux, T. Mahmoud, C. Collard, M.-C. Lemaire, and V. Litvin, "TeV electron and photon saturation studies," *CMS Note* **2006/004** (2006).
- [739] D. Acosta et al., "Measurement of the Cross Section for Prompt Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV," *Phys. Rev. Lett.* **95** (2005) 022003.
doi:10.1103/PhysRevLett.95.022003.
- [740] J. Weng et al., "Search for ADD Direct Graviton Emission in Photon plus Missing Transverse Energy Final States at CMS," *CMS Note* **2006/129** (2006).
- [741] R. C. Myers and M. J. Perry, "Black holes in higher dimensional space-times," *Ann. Phys.* **172** (1986) 304.
- [742] S. W. Hawking, "Particle creation by black holes," *Commun. Math. Phys.* **43** (1975) 199–220.
- [743] P. C. Argyres, S. Dimopoulos, and J. March-Russell, "Black holes and sub-millimeter dimensions," *Phys. Lett.* **B441** (1998) 96–104, arXiv:hep-th/9808138.
- [744] C. M. Harris, P. Richardson, and B. R. Webber, "CHARYBDIS: A black hole event generator," *JHEP* **08** (2003) 033, arXiv:hep-ph/0307305.
- [745] K. D. Lane, "Technicolor 2000," arXiv:hep-ph/0007304.
- [746] P. Kreuzer, "Search for Technicolour at CMS in the $\rho_{TC} \rightarrow WZ$ channel," *CMS Note* (2006/135).
- [747] B. Betev, D. Bourilkov, and S. S. Mavrodiev, "Structure Functions of Pion and Nucleon Determined from High Mass Muon Pair Production." JINR-E2-85-312.
- [748] Super-Kamiokande Collaboration, Y. Fukuda et al., "Evidence for oscillation of atmospheric neutrinos," *Phys. Rev. Lett.* **81** (1998) 1562–1567,
arXiv:hep-ex/9807003.
- [749] S. Esen and R. Harris, "CMS Sensitivity to Quark Contact Interactions using Dijets," *CMS Note* **2006/071** (2006).
- [750] J. C. Pati and A. Salam, "Anomalous lepton-hadron interactions and gauge models," *Phys. Rev.* **D11** (1975) 1137–1154. doi:10.1103/PhysRevD.11.1137.

- [751] CMS Collaboration, N. Gninenko, M. Kirsanov, N. Krasnikov, and N. Matveev, "Detection of Heavy Majorana Neutrinos and Right-Handed Bosons," *CMS Note* **2006/098** (2006).
- [752] Particle Data Group Collaboration, K. Hagiwara et al., "Review of particle physics," *Phys. Rev.* **D66** (2002) 010001.
- [753] G. Barenboim, J. Bernabeu, J. Prades, and M. Raidal, "Constraints on the W(R) mass and CP-violation in left-right models," *Phys. Rev.* **D55** (1997) 4213–4221, arXiv:hep-ph/9611347.
- [754] DØ Collaboration, S. Abachi et al., "Search for Right-Handed W Bosons and Heavy W' in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev. Lett.* **76** (1996) 3271–3276, arXiv:hep-ex/9512007.
- [755] A. Datta, M. Guchait, and D. P. Roy, "Prospect of heavy right-handed neutrino search at SSC / LHC energies," *Phys. Rev.* **D47** (1993) 961–966, arXiv:hep-ph/9208228.
- [756] CMS Collaboration, K. Karafasoulis, A. Kyriakis, H. Petrakou, and K. Mazumdar, "Little Higgs Model and Top-like Heavy Quark at CMS," *CMS Note* **2006/079** (2006).
- [757] J. D'Hondt, S. Lowette, G. Hammad, J. Heyninck, and P. Van Mulders, "Observability of same-charge lepton topology in di-leptonic $t\bar{t}$ events," *CMS Note* **2006/065** (2006).
- [758] F. Larios and F. Penunuri, "FCNC production of same sign top quark pairs at the LHC," *J. Phys.* **G30** (2004) 895–904, arXiv:hep-ph/0311056.
- [759] Y. P. Gouz and S. R. Slabospitsky, "Double top production at hadronic colliders," *Phys. Lett.* **B457** (1999) 177–185. doi:10.1016/S0370-2693(99)00516-X.
- [760] C.-X. Yue, Z.-J. Zong, L.-L. Xu, and J.-X. Chen, "Associated production of the top-pions and single top at hadron colliders," *Phys. Rev.* **D73** (2006) 015006, arXiv:hep-ph/0601058.
- [761] S. Kraml and A. R. Raklev, "Same-sign top quarks as signature of light stops at the LHC," *Phys. Rev.* **D73** (2006) 075002, arXiv:hep-ph/0512284.
- [762] S. Bitjukov and N. Krasnikov, "On the observability of a signal above background," *Nucl. Instrum. and Methods* **A452** (2000) 518–524. doi:10.1016/S0168-9002(00)00454-X.
- [763] E. Lehmann, "A Theory of some Multiple Decision Problems," *The Annals of Mathematical Statistics* **28** 1 (1957).
- [764] R. O'Neill and G. Wetherill, "The Present State of Multiple Comparison Methods," *Journal of the Royal Statistical Society, Series B (Methodological)* **33** 2 (1971).
- [765] J. Hauser, "Search for New Physics in Tails of Distributions," 2006. Private communications, paper in preparation.
- [766] DØ Collaboration, V. M. Abazov et al., "Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using kinematic characteristics of lepton + jets events," *Phys. Lett.* **B626** (2005) 45–54, arXiv:hep-ex/0504043. doi:10.1016/j.physletb.2005.08.104.

- [767] B. Abbott et al., "Search for New Physics in $e\mu X$ Data at DØ Using Sleuth: A Quasi-Model-Independent Search Strategy for New Physics," *Phys. Rev.* **D62** (2000). doi:10.1103/PhysRevD.62.092004.
- [768] B. Abbott et al., "A Quasi-Model-Independent Search for New Physics at Large Transverse Momentum," *Phys. Rev.* **D64** (2001) arXiv:hep-ex/0011067.
- [769] C. Finley et al., "On the evidence for clustering in the arrival directions of AGASA's ultrahigh energy cosmic rays," *Astroparticle Physics* **21** (2004).
- [770] S. Frixione, P. Nason, and B. R. Webber, "Matching NLO QCD and parton showers in heavy flavour production," *JHEP* **08** (2003) 007, arXiv:hep-ph/0305252.
- [771] L. Lonnblad, "Correcting the colour-dipole cascade model with fixed order matrix elements," *JHEP* **05** (2002) 046, arXiv:hep-ph/0112284.
- [772] S. Frixione and B. R. Webber, "Matching NLO QCD computations and parton shower simulations," *JHEP* **06** (2002) 029, arXiv:hep-ph/0204244.
- [773] F. Krauss, "Matrix elements and parton showers in hadronic interactions," *JHEP* **08** (2002) 015, arXiv:hep-ph/0205283.
- [774] S. Mrenna and P. Richardson, "Matching matrix elements and parton showers with HERWIG and PYTHIA," *JHEP* **05** (2004) 040, arXiv:hep-ph/0312274.
- [775] J. Butterworth, S. Butterworth, B. Waugh, W. Stirling, and M. Whalley, "The CEDAR project," arXiv:hep-ph/0412139.
- [776] LEP Collaboration, "A combination of preliminary electroweak measurements and constraints on the standard model," arXiv:hep-ex/0412015.
- [777] D. R. Stump, "A new generation of CTEQ parton distribution functions with uncertainty analysis." Prepared for 31st International Conference on High Energy Physics (ICHEP 2002), Amsterdam, The Netherlands, 24-31 Jul 2002.
- [778] W. J. Stirling, A. D. Martin, R. G. Roberts, and R. S. Thorne, "MRST parton distributions," *AIP Conf. Proc.* **747** (2005) 16–21.
- [779] M. Botje, "A QCD analysis of HERA and fixed target structure function data," *Eur. Phys. J.* **C14** (2000) 285–297, arXiv:hep-ph/9912439.
- [780] S. I. Alekhin, "Global fit to the charged leptons DIS data: $\alpha(s)$, parton distributions, and high twists," *Phys. Rev.* **D63** (2001) 094022, arXiv:hep-ph/0011002.
- [781] G. Altarelli and G. Parisi, "Asymptotic Freedom in Parton Language," *Nucl. Phys.* **B126** (1977) 298.
- [782] L. Lonnblad, "ARIADNE version 4: A Program for simulation of QCD cascades implementing the color dipole model," *Comput. Phys. Commun.* **71** (1992) 15–31.
- [783] CDF Collaboration, F. Abe et al., "Evidence for color coherence in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Rev.* **D50** (1994) 5562–5579. doi:10.1103/PhysRevD.50.5562.

- [784] **DØ** Collaboration, B. Abbott et al., "Color coherent radiation in multijet events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV," *Phys. Lett.* **B414** (1997) 419–427.
doi:10.1016/S0370-2693(97)01190-8.
- [785] I. G. Knowles et al., "QCD Event Generators," arXiv:hep-ph/9601212.
- [786] **ALEPH** Collaboration, D. Buskulic et al., "Properties of hadronic Z decays and test of QCD generators," *Z. Phys.* **C55** (1992) 209–234.
- [787] **DELPHI** Collaboration, P. Abreu et al., "Tuning and test of fragmentation models based on identified particles and precision event shape data," *Z. Phys.* **C73** (1996) 11–60. doi:10.1007/s002880050295.
- [788] **OPAL** Collaboration, G. Alexander et al., "A Comparison of b and (u d s) quark jets to gluon jets," *Z. Phys.* **C69** (1996) 543–560.
- [789] B. Andersson, G. Gustafson, and B. Soderberg, "A GENERAL MODEL FOR JET FRAGMENTATION," *Z. Phys.* **C20** (1983) 317.
- [790] **ALEPH** Collaboration, A. Heister et al., "Study of the fragmentation of b quarks into B mesons at the Z peak," *Phys. Lett.* **B512** (2001) 30–48, arXiv:hep-ex/0106051.
- [791] **OPAL** Collaboration, G. Abbiendi et al., "Inclusive analysis of the b quark fragmentation function in Z decays at LEP. ((B))," *Eur. Phys. J.* **C29** (2003) 463–478, arXiv:hep-ex/0210031.
- [792] **SLD** Collaboration, K. Abe et al., "Measurement of the b-quark fragmentation function in Z^0 decays," *Phys. Rev.* **D65** (2002) 092006.
doi:10.1103/PhysRevD.65.092006.
- [793] M. G. Bowler, "e+ e- Production of Heavy Quarks in the String Model," *Zeit. Phys.* **C11** (1981) 169.
- [794] C. Peterson, D. Schlatter, I. Schmitt, and P. M. Zerwas, "Scaling Violations in Inclusive e^+e^- Annihilation Spectra," *Phys. Rev.* **D27** (1983) 105.
- [795] V. Kartvelishvili, A. Likhoded, and V. Petrov, "On the fragmentation functions of heavy quarks into hadrons," *Phys. Lett.* **B78** (1978) 615.
- [796] E. Norrbin and T. Sjostrand, "Production and hadronization of heavy quarks," *Eur. Phys. J.* **C17** (2000) 137–161, arXiv:hep-ph/0005110.
- [797] R. Engel and J. Ranft, "Color singlet exchange between jets and the PHOJET Monte Carlo," *Nucl. Phys. Proc. Suppl.* **75A** (1999) 272–274.
doi:10.1016/S0920-5632(99)00263-7.
- [798] **CDF** Collaboration, T. Affolder et al., "Charged jet evolution and the underlying event in proton anti-proton collisions at 1.8 TeV," *Phys. Rev.* **D65** (2002) 092009.
doi:10.1103/PhysRevD.65.092009.
- [799] P. Nason et al., "Bottom production," arXiv:hep-ph/0003142.
- [800] T. Sjostrand and P. Z. Skands, "Transverse-momentum-ordered showers and interleaved multiple interactions," *Eur. Phys. J.* **C39** (2005) 129–154,
arXiv:hep-ph/0408302. doi:10.1140/epjc/s2004-02084-y.

- [801] S. Sapeta and K. Golec-Biernat, "Total, elastic and diffractive cross sections at LHC in the Miettinen-Pumplin model," *Phys. Lett.* **B613** (2005) 154–161, arXiv:hep-ph/0502229.
- [802] D. J. Lange, "The EvtGen particle decay simulation package," *Nucl. Instrum. Meth.* **A462** (2001) 152–155.
- [803] P. Bartalini, "Supporting Monte Carlo Generators at the LHC," in *Proceedings of the 10th International Conference on B-Physics at Hadron Machines*, M. Biasini and S. Erhan, eds. Assisi (Perugia), Italy, June, 2005. Published in *Nucl. Phys B - Proceedings Supplements*, Vol 156, 1.
- [804] Z. Sullivan, "Fully differential W' production and decay at next-to-leading order in QCD," *Phys. Rev.* **D66** (2002) 075011, arXiv:hep-ph/0207290.
- [805] M. A. Dobbs et al., "Les Houches guidebook to Monte Carlo generators for hadron collider physics," arXiv:hep-ph/0403045.
- [806] "HIJING Web Site." Located at <http://www-nsdth.lbl.gov/~xnwang/hijing/index.html>.
- [807] T. Gleisberg et al., "Event generator for the LHC," arXiv:hep-ph/0508315.
- [808] E. Boos et al., "Generic user process interface for event generators," arXiv:hep-ph/0109068.
- [809] I. P. Lokhtin and A. M. Snigirev, "Nuclear geometry of jet quenching," *Eur. Phys. J.* **C16** (2000) 527–536, arXiv:hep-ph/0004176. doi:10.1007/s100520000437.
- [810] I. P. Lokhtin and A. M. Snigirev, "Fast simulation of jet quenching in ultrarelativistic heavy ion collisions," arXiv:hep-ph/0406038. SINP MSU-13/752.
- [811] I. P. Lokhtin and A. M. Snigirev, "Fast simulation of flow effects in central and semi-central heavy ion collisions at LHC," arXiv:hep-ph/0312204.
- [812] CDF Collaboration, R. D. Field, "The underlying event in hard scattering processes," *eConf* **C010630** (2001) P501, arXiv:hep-ph/0201192.
- [813] CompHEP Collaboration, E. Boos, V. Bunichev, M. Dubinin, L. Dudko, V. Edneral, V. Ilyin, A. Kryukov, V. Savrin, A. Semenov, and A. Sherstnev, 2004. Web site at <http://theory.sinp.msu.ru/comphep>. See also [354].
- [814] I. P. Lokhtin, L. I. Sarycheva, and A. M. Snigirev, "The method for analysing jet azimuthal anisotropy in ultrarelativistic heavy ion collisions," *Phys. Lett.* **B537** (2002) 261–267, arXiv:hep-ph/0203144. doi:10.1016/S0370-2693(02)01913-5.
- [815] M. Carena, A. Daleo, B. A. Dobrescu, and T. M. P. Tait, " Z' gauge bosons at the Tevatron," *Phys. Rev.* **D70** (2004) 093009. doi:10.1103/PhysRevD.70.093009.
- [816] C. Balazs, J.-w. Qiu, and C. P. Yuan, "Effects of QCD resummation on distributions of leptons from the decay of electroweak vector bosons," *Phys. Lett.* **B355** (1995) 548–554. doi:10.1016/0370-2693(95)00726-2.
- [817] C. Balazs and C. P. Yuan, "Soft gluon effects on lepton pairs at hadron colliders," *Phys. Rev.* **D56** (1997) 5558–5583. doi:10.1103/PhysRevD.56.5558.

- [818] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, "Uncertainties of predictions from parton distributions. I: Experimental errors," *Eur. Phys. J.* **C28** (2003) 455–473, arXiv:hep-ph/0211080.
- [819] J. H. Holland, "Adaptation in natural and artificial systems". The University of Michigan Press, Ann Arbor, 1975.
- [820] D. E. Goldberg, "Genetic algorithms in search, optimization and machine learning". Addison Wesley, 1989.
- [821] S. Abdullin, "Genetic Algorithm for SUSY Trigger Optimization In CMS Detector At LHC," *NIM A* **502** (2003) 693–695. doi:10.1016/S0168-9002(03)00546-1.
- [822] TOTEM Collaboration, "TOTEM Technical Design Report," *CERN/LHCC 2004-002* (2004).
- [823] TOTEM Collaboration, "Addendum to the TOTEM-TDR," *CERN/LHCC 2004-020* (2004).
- [824] V. Avati and K. Österberg, "TOTEM forward measurements: leading proton acceptance," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4avati.pdf>.
- [825] J. Kalliopuska et al., "TOTEM forward measurements: exclusive central diffraction," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at http://www.desy.de/~heralhc/proceedings/wg4pXp_heralhc.pdf.
- [826] "The MAD-X Program, Methodical Accelerator Design." Information available at <http://www.cern.ch/mad>.
- [827] M. Arneodo et al., "Diffractive Higgs: CMS/TOTEM Level-1 Trigger Studies," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg4arneodo.pdf>.
- [828] R. Croft, "(In preparation)". PhD thesis, University of Bristol, 2006.
- [829] F. Oljemark, "First level triggering of diffractively produced low-mass Higgs at The Large Hadron Collider," Master's thesis, University of Helsinki, 2006.
- [830] G. Bruni et al., "Leading proton production in ep and pp , experiments: how well do high-energy physics Monte Carlos reproduce the data?," in *Proceedings of the HERA-LHC Workshop*. CERN/DESY, January, 2005. Available at <http://www.desy.de/~heralhc/proceedings/wg5lps.pdf>.
- [831] F. Ferro, "Diffractive Higgs in CMS/TOTEM: study of L1 trigger conditions from T1 and T2," *TOTEM Note 04-2005* (2005).
- [832] B. Cox and J. Forshaw, "Herwig for diffractive interactions," *Comput. Phys. Commun.* **144** (2002) 104–110. doi:10.1016/S0010-4655(01)00467-2.
- [833] J. C. Collins, "Light-cone variables, rapidity and all that," arXiv:hep-ph/9705393.
- [834] M. G. Albrow and A. Rostovtsev, "Searching for the Higgs at hadron colliders using the missing mass method," arXiv:hep-ph/0009336.

-
- [835] M. G. Albrow, "Double pomeron physics at the LHC," *AIP Conf. Proc.* **792** (2005) 509–514, [arXiv:hep-ex/0507095](https://arxiv.org/abs/hep-ex/0507095).

